

REVIEW ARTICLE

Open access

# Systematics and distribution of the true monitors *Varanus* Merrem, 1820 (Sauria, Varanidae) on the Iranian Plateau

Narges Ghanbarinia<sup>ID</sup>, Rasoul Karamiani<sup>ID\*</sup>, Ahmad Gharezi<sup>ID</sup>, & Nasrullah Rastegar-Pouyani<sup>ID</sup>

Department of Biology, Faculty of Science, Razi University, 6714967346 Kermanshah, Iran.

## Abstract

Varanoidea is an ancient group of anguimorph lizards, the anguimorph genus *Varanus* includes today's monitor lizards, a clade of mid-sized to large, squamates widely distributed in tropical and subtropical regions. biogeographic analysis suggests an origin of varaniform lizards in East Asia. According to most studies, monitors grouped in three clades concordant with their geographic distribution: African, Indo-Asian, and Indo-Australian. There are currently 88 valid species assigned to *Varanus* that divided into 11 subgenera. In the Iranian Plateau, there are four species of monitors from two subgenera *Psammosaurus* including *Varanus griseus*, *Varanus caspius*, *Varanus nesterovi* and *Empagusia* include the species *Varanus bengalensis*. *V. caspius* is mostly present in the central parts of Iran, two species *V. griseus* and *V. nesterovi* being distributed from Kermanshah province to Shiraz and also in Iraq. The end of distribution of the species *V. bengalensis* is in the southeast of Iran including Sistan-Baluchestan and Hormozgan Provinces.

**Key words:** *Varanus*, monitor, Varanidae, Distribution, Iranian plateau, Asia.

Received:  
26 February 2025

Accepted:  
23 May 2025

## INTRODUCTION

Varanoidea is an ancient group of anguimorph lizards, comprising two extant New World helodermatids (*Heloderma horridum* and *Heloderma suspectum*), the Bornean earless monitor (*Lanthanotus borneensis*), and the Old-World monitors (*Varanus*) (Ast, 2001). The anguimorph genus *Varanus* encompasses today's monitor lizards (Varanidae Hardwicke & Gray, 1827), a clade of mid-sized to large, mostly carnivorous (though some are frugivorous) squamates widely distributed in tropical and subtropical regions (Fig. 1; Ivanov et al., 2017). In animals, body size dramatically affects life-history traits and ecology, and it is the most commonly used measurement in macroevolutionary studies (Wilson, 1975). Among terrestrial vertebrates, monitor lizards (*Varanus*) exhibit the greatest variation in body size within a single terrestrial vertebrate genus (Pianka, 1995). Extant monitors include island giants like the Komodo dragon (*V. komodoensis*) (up to 3 m long and 100 kg) and desert dwarves like the short-tailed goanna (*V. brevicauda*) (0.2 m and 0.016 kg), which vary by orders of magnitude (Brennan et al., 2021). Despite a conservative body plan, monitor lizards in Africa, Asia, and Australia are ecologically diverse. They also show specific ecological and morphological adaptations and can be found at home in trees, among rocks, in burrows, and swimming through watercourses and even the open ocean (Pianka, 1995; Bucklitsch et al., 2016). Monitor lizards inhabit a variety of environments, from deserts to rainforests (Bucklitsch et al., 2016). Most varanids are moderate to large in size with thick skin containing numerous rows of small, rounded scales circling the body. Monitors have well-developed limbs; their tail is long to very long and lacks caudal autotomy (Vitt & Caldwell, 2013). The greatest morphological diversity is

\*Corresponding Author: ✉ : [r.karamiani@razi.ac.ir](mailto:r.karamiani@razi.ac.ir)

## HOW TO CITE THIS ARTICLE:


Ghanbarinia, N., Karamiani, R., Gharzi, A., Rastegar-Pouyani, N. (2025). 'Systematics and distribution of the true monitors *Varanus* Merrem, 1820 (Sauria, Varanidae) on the Iranian Plateau', *Iranian Journal of Animal Biosystematics*, 21(1), pp. 75-91. doi: 10.22067/ijab.2025.92359.1094





Currently, no extant monitor lizard species are known from European territory, as geographically defined by an eastern border following the Ural Mountains and the Ural River, through the Caspian Sea to the eastern end of the main chain of the Great Caucasus, from which it runs westwards to the Black Sea. From there, it continues through the Bosphorus, the Marmara Sea, and the Dardanelles towards the Aegean Sea, where the islands on the Anatolian shelf are excluded (Mertens and Wermuth, 1960). However, fossil representatives of this family were present in Europe until the Middle Pleistocene (Georgalis et al., 2017; Böhme et al., 2023). The fossil record of Varanidae, although still very incomplete, demonstrates great diversity across most of the Cenozoic and a wide geographic distribution. This encompasses broad areas that are currently well outside the range of extant forms, such as North America and Europe (Georgalis et al., 2021; Molnar, 2004). In fact, the varanid fossil record is far richer in Europe than on any other continent, spanning from the earliest Eocene up to the Middle Pleistocene (Augé, 2005; Augé et al., 2022; Fejérváry, 1935; Georgalis et al., 2017; Hofstetter, 1969; Villa et al., 2018; Georgalis et al., 2023). Most studies use these fossils and molecular data to determine interspecific relationships (Arida & Böhme, 2010). Several varanid paleospecies have been described from European and neighboring fossil sites. Those described before Estes (1983) were already synonymized by this author with *V. marathonsensis* (Villa et al., 2018). The latter authors recognized only two European paleospecies, namely *V. marathonsensis* and *V. mokrensis*. This seems problematic as long as only osteological characters are available (Böhme et al., 2023). The mitochondrial phylogeny by Ast (2001) is probably the most comprehensive and reliable study inferring relationships among extant varanid lizards because it is well sampled across taxonomic classification and geographic distribution (Arida & Böhme, 2010). In this phylogeny, extant species are grouped into three clades concordant with their geographic distribution: African, Indo-Asian, and Indo-Australian (Villa et al., 2018).

Given the phylogenetic hypothesis of Conrad (2012), no fewer than eight major *Varanus* lineages must have been present. These radiations include groups of Indo-Asian A monitors (the *Empagusia*, *Indo-Varanus*, and *Tecto-Varanus* clades), the *Dendro-Varanus*-*Soteriosaurus* clade, the Australasian clade, the Indo-Asian B clade, the lineage including *Varanus griseus*, and the *Polydaedalus* clade (Conrad et al., 2012). Relationships among anguimorph lizard groups have been contentious, particularly with regard to the placement of fossil taxa (Conrad, 2008; Conrad et al., 2011; Pyron, 2017). Analysis of morphological data in concert with novel phylogenomic data are largely consistent with previous assessments; however, Brennan et al. (2021) provide new insights into the phylogenetics of living members of *Varanus*. This phylogeny of *Varanus* highlights the adaptive capacity of these amazing lizards, emphasizing their ability to fill available niches. Overall, Brennan et al. (2021) suggest a more recent timeline for the diversification of modern varanid lizards when compared to other phylogenetic studies, with a crown age in the early-to-mid Oligocene. This timing suggests *Varanus* potentially dispersed into the Indo-Australian region shortly after the collision of the Australian and Asian plates. If this is true, the connection of Sahul to Sundaland likely facilitated the dispersal of monitor lizards across an Indonesian island bridge, and extensive overwater dispersal seems less probable. This pattern is consistent with the adaptive radiation of Australopapuan elapid snakes (Keogh, 1998) and python snakes (Reynolds et al., 2014; Esquerre et al., 2019), from Asian origins (Fig. 1; Brennan et al., 2021). The most recent study on the phylogenetic relationships of varanids was carried out by Georgalis et al. (2023) after the discovery of two fossil teeth from the Early Miocene of Hüenerbach, Switzerland. In this study, to obtain the optimal tree, they performed parsimony-based character optimization on denticles based on the phylogeny of Pyron et al. (2013) (Fig. 2; Georgalis et al., 2023). Levshakova (1986) and Stanner (2004) considered *Psammosaurus* to be composed of three species, namely the extant *V. griseus* and the two fossil taxa *V. darevskii* and *V. marathonsensis*. Ivanov et al. (2018), as well as Villa et al. (2018), in contrast, assigned the last-named species to the Indian clade where *V. bengalensis* is also accommodated (the *Empagusia* group of *Varanus*). *Varanus darevskii* from the Pliocene of Tadzhikistan, however, is still regarded as a close relative or even predecessor of the extant desert monitors, and, particularly, on biogeographical grounds, of *V. caspius*. Thus, the SW Asian origin of the *Psammosaurus* clade is made plausible by



**FIGURE 2.** Parsimony-based character optimization on the tooth serrations based on the phylogeny of Pyron et al. (2013) (Georgalis et al., 2023)

*V. darevskii*, the extinct predecessor and Pliocene ancestor of *V. caspius* (Levchakova, 1986) from Tadzhikistan. Apart from the distinct morphological differences between the nominal taxa *V. griseus*, *V. nesterovi*, and *V. caspius* (shape of nostril, neck scalation, dorsal color pattern, tail shape, and tail coloration), the chorological pattern of the three parapatric forms also supports their full specific rather than subspecific status (Böhme et al., 2023).

#### SYSTEMATICS ACCOUNT

##### Order Squamata Oppel, 1811

##### Superfamily Varanoidea Camp, 1923

##### Family Varanidae Gray, 1827

##### Genus *Varanus* Merrem, 1820

There are currently 88 valid species assigned to *Varanus* (Uetz et al., 2024). This genus is divided into 11 subgenera, eight of which are represented in the Pacific East. Of these, one (*Philippinosaurus*) occurs only in the Philippines, one (*Soterosaurus*) is primarily found west of Weber's Line, and five (*Euprepiosaurus*, *Hapterosaurus*, *Odatia*, *Papusa*, *Varanus*) are restricted to Australia, New Guinea, and some of the Lesser Sunda Islands. The monotypic subgenus *Solomonsaurus* is restricted to parts of the western Solomon Islands (McCoy, 2015; Weijola, 2017; Uetz et al., 2024). Other subgenera include *Psammosaurus*, *Polydaedalus*, and *Empagusia* (Böhme, 2003).

##### Subgenus *Soterosaurus* Ziegler & Böhme, 1997

**Distribution:** From Sri Lanka through continental and insular Southeast Asia including the Philippines as well as the Greater and Lesser Sunda Islands, Sulawesi, and the northern Moluccas (Koch et al., 2013).

**Subgenus *Varanus* Merrem, 1820**

**Distribution:** Australia, southern part of New Guinea, and some Lesser Sunda Islands (Koch et al., 2013).

**Subgenus *Papusaurus* Mertens, 1962**

**Distribution:** Restricted to New Guinea.

The monotypic subgenus *Papusaurus* contains only one species, *V. salvadorii* (Koch et al., 2013).

**Subgenus *Odatia* Gray, 1838**

**Distribution:** Australia, New Guinea, Timor and off-shore islands.

The subgenus *Odatia* of monitor lizards underwent a major radiation on the Australian. three members of this subgenus, i.e., *V. auffenbergi*, *V. similis*, and *V. timorensis*, inhabit islands of the IndoAustralian Archipelago (Koch et al., 2013).

**Subgenus *Philippinosaurus* MERTENS, 1959**

**Distribution:** Restricted to the Philippine islands of Luzon and Panay.

The subgenus *Philippinosaurus* comprises the only fructivorous monitor lizard species that feed mainly on leaves and fruits. All species are endemic to the Philippines (Koch et al., 2013).

**Subgenus *Euprepiosaurus* Fitzinger, 1843**

**Distribution:** Moluccas, New Guinea, Northern Australia, and various Pacific island groups such as the Bismarck Archipelago, the Solomons, and the Marshall Islands.

The subgenus *Euprepiosaurus* comprises two monophyletic clades, viz. the species groups of Pacific or Mangrove Monitors around *V. indicus* and the Tree Monitors around *V. prasinus* (Koch et al., 2013).

**Subgenus *Haptesaurus* Buckitsch, Bohme & Koch, 2016**

**Distribution.** Members of the subgenus *Hapturosaurus* are restricted to the tropical rainforests of New Guinea and several offshore islands, such as Waigeo, Batanta, and the Aru Islands, as well as Northern Australia. During Pleistocene sea level depressions of up to 140 meters below present levels, all these island regions belonging to the Sahul Shelf were interconnected by subaerial land bridges (Voris, 2000; Bucklitsch et al., 2016).

*Hapturosaurus* is a monophyletic group (Ast, 2001; Ziegler et al., 2007) of medium-sized arboreal lizards (Collar et al., 2011) from New Guinea and northern Australia. This group was formerly placed as a species group within the subgenus *Euprepiosaurus*. The name *Hapturosaurus* refers to the autapomorphy of the prehensile tail found in the highly arboreal tree monitors of the *V. prasinus* species group (Bucklitsch et al., 2016).

**Subgenus *Solomonsaurus* Buckitsch, Bohme & Koch, 2016**

**Distribution.** The subgenus *Solomonsaurus* is endemic to the Solomon Islands southeast of New Guinea. it is known only from the islands of San Jorge (St. George), Santa Isabel, and Bougainville (Böhme & Ziegler 2007).

Originally, Mertens (1941) described *Varanus spinulosus* as a subspecies of the widespread *Varanus indicus*, based on his multiple species concept of Pacific monitors (Mertens, 1942). Later, the taxon was elevated to full species status by Sprackland (1994) and was initially placed in the subgenus *Euprepiosaurus* by Ziegler & Böhme (1997). However, *V. spinulosus* recently had to be removed from this subgenus because it lacks the unique synapomorphies of this monitor in genital morphology (Böhme & Ziegler, 2007). Therefore, the systematic position of this monitor lizard species was considered *incertae sedis* (Koch et al., 2010) and required review. In 2016, it was placed in the new subgenus *Solomonsaurus*. The name *Solomonsaurus* refers to the Solomon Islands, which is the distribution range of the only recognized species within this currently monotypic subgenus (Bucklitsch et al., 2016).

**Subgenus *Polydaedalus* Wagler, 1830**

**Distribution:** the African steppe monitors (Bohme & Ziegler, 1997).

The species of *Polydaedalus* mostly inhabit savannah habitats in Africa (Bucklitsch et al., 2016).

**Subgenus *Empagusia* Gray, 1838**

**Distribution:** From southeastern Iran and Afghanistan through Pakistan, India, Bangladesh and Nepal to Myanmar, Thailand, Malaysia, and the Greater Sunda Islands, Indonesia.

The subgenus *Empagusia* represents a paraphyletic clade (Koch et al., 2013).

**Subgenus *Psammosaurus* Fitzinger, 1826**

**Distribution:** northern Africa, Arabia, Middle East, Central Asia and southern Asia.

Species of the monotypic subgenus *Psammosaurus* are adapted to arid environments in Africa, Arabia, and southern Asia. This taxon is sister to the *Polydaedalus* clade in recent molecular phylogenies (Ast, 2001; Collar et al., 2011).

**Manitors in the Iranian plateau**

*Varanus psammosaurus griseus* (Daudin, 1803)

*Tupinambis griseus* Daudin 1803 (Type locality: Dardsha, coast of the Caspian Sea)

*Varanus scincus* Merrem 1820

*Tupinambis arenarius* Geoffroy 1827

*Varanus arenarius* Duméril & Bibron 1836

*Monitor terrestris* Calori 1857

*Psammosaurus terrestris* Rabl Rückhard 1881

*Psammosaurus scincus* Blanford 1881

*Varanus griseus* Boulenger 1885

*Psammosaurus griseus* Sixta 1900

*Varanus (Psammosaurus) griseus* Mertens 1942

*Varanus griseus griseus* Gardner 2013

*Varanus griseus* Šmíd et al. 2014

*Varanus (Psammosaurus) griseus* Bucklitsch et al. 2016

*Varanus griseus* Bar et al. 2021

*Varanus griseus* Böhme et al. 2023

The **desert monitor lizard** has a concave head and a nostril in the form of a diagonal slit that narrows anteriorly and is closer to the eye. It possesses relatively weakly enlarged glands on the neck, a dorsal side usually with dark and/or distinct cross-stripes, and lighter spots on the ground that are darker. Its tail is round in cross-section with distinct cross-stripes along its length (Fig. 3; Böhme, 2015). *Varanus griseus* (Daudin, 1803) is a terrestrial species with a desert-sandy distribution pattern whose population includes two subspecies: *V. g. griseus* in North Africa, Arabia, and the Near East, and *V. g. koniecznyi* Mertens, 1954 in southern Pakistan and northwestern India (Mertens, 1942, 1954; Böhme, 1997, 2003, 2015). The gray desert monitor is found in the Provinces of Khuzestan, Ilam, Kermanshah, Bushehr, and Fars (Fig. 7; Safaei Mahro et al., 2015).

The zoogeographical origin of Desert Monitor is Saharo-Turano-Sindian region (Anderson, 1999; Baig et al., 2008; Bauer et al., 2017; Bayless, 2002; Damadi et al., 2017; Escoriza et al., 2011; Jablonski et al., 2021; Khan, 2004; Mertens, 1954; Malakhov & Chirikova, 2018; Sindaco et al., 2000; Akman et al., 2023).

***Varanus psammosaurus caspius* (Eichwald, 1831)**

*Psammosaurus caspius* Eichwald 1831 (Type locality: Dardsha Peninsula, southeastern coast of Caspian Sea, Turkmenistan).





**FIGURE 3.** *Varanus griseus*. The sample was found in the south of Dehloran city, Ilam province (Fathnia et al., 2009).

*Varanus caspius* Gray 1845

*Varanus griseus caspius* Anderson 1963

*Varanus griseus caspius* Nilson & Andren 1981

*Varanus griseus caspius* Khan 2003

*Varanus griseus caspius* Heidari et al. 2010

*Varanus caspius* Böhme et al. 2023

In 2023, Boehm et al. declared the *V. caspius* a separate species from the *V. griseus*, based on morphological differences, habitat, and distribution of Iranian plateau monitors and the results of the Brennan et al (2021) study (Fig. 4).

The following are some of the characteristics of the scaly fish: the anterior half of the tail is more or less circular in transverse section, narrow in the posterior half, compressed on both sides in transverse section, and has a distinct blade on the upper surface; the ventral scales are in 110-125 transverse rows from the collar to the groin; the transverse stripes on the dorsal area are red-brown; and the tail has 13-19 dark transverse stripes; the end of the tail (usually the distal third) is uniformly yellowish and patternless (Rastegar-Pouyani et al., 2007).

*V. g. caspius* (Eichwald, 1831) has a vast distribution from central Iran (Qom) and the eastern coast of the Caspian Sea in the west to Tadzhikistan in the east (Fig. 7; Ehrlich, 2013; Chirikova et al., 2019; Böhme, 2023).

***Varanus empagusia bengalensis*** (Daudin, 1802)

*Tupinambis bengalensis* Daudin 1802

*Tupinambis cepedianus* Daudin 1802

*Lacerta argus* Daudin 1802

*Varanus punctatus* Merrem 1820

*Varanus taraguira* Merrem 1820

*Varanus bengalensis* Duméril & Bibron 1836

*Monitor inornatus* Schlegel 1839

*Monitor cepedianus* Schlegel 1839

*Uvaranus lunatus* Gray 1845

*Varanus dracaena* Theobald 1868



**FIGURE 4.** *Varanus caspius*. The Zoological Museum specimen of Razi University has been used for photographing.

*Varanus lunatus* Anderson 1871

*Varanus dracaena* Anderson 1871

*Varanus bengalensis* Mertens 1942

*Varanus (Empagusia) bengalensis* Koch et al. 2013

*Varanus (Empagusia) bengalensis* Bucklitsch et al. 2016

The Bengal Monitor (*Varanus bengalensis*) can reach a snout-vent length (SVL) of around 180 cm, with a 100 cm tail. It's terrestrial in habit, though young individuals can be arboreal (Fig. 5; Kabir et al., 2009). This species is primarily omnivorous; however, scavenging is frequently observed. It's found in all habitats throughout Bangladesh. As a large and ecologically flexible species, *Varanus bengalensis* is commonly encountered in many ecological settings, including human-modified habitats. Its global distribution ranges from southeastern Iran to southern Asia (Jaman et al., 2024). In Iran, the Bengal Monitor, also sometimes called the Bengal Tiger, is present in the Provinces of Sistan and Baluchestan, Kerman, and Hormozgan (Fig. 7; Safaei-Mahroo et al., 2015).

*Varanus psammosaurus nesterovi* Böhme, Ehrlich, Milto, Orlov & Scholz, 2015

*Varanus nesterovi* Böhme, Ehrlich, Milto, Orlov & Scholz 2015 (Type locality: "Biare," currently Byara Village, Muhafazat as Sulaymaniyah, Iraq (35°13'50" N 46°07'15" E), 1086 m elevation)

*Varanus griseus* Tuck 1971

*Varanus griseus caspius* Mertens 1973

*Varanus griseus caspius* Fathinia et al. 2009

*Varanus (Psammosaurus) nesterovi* Bucklitsch et al. 2016

*Varanus (Psammosaurus) nesterovi* Böhme et al. 2023

Nesterov's desert monitor A large growing representative of the section *Psammosaurus Varanus* (total length up to 120 cm) easily distinguished from *V. griseus*, 1.its broad, short head with a convex profile. 2. nostrils round to vertically oval, not quite in front of the eyes. 3.scaly neck of strongly enlarged scales and spines, strongest on the sides of the neck, which may even form curved spines. 4.dorsal side uniformly sandy, without well-visible dark cross-stripes, at best with only rudimentary and indistinct streaks of a dark pattern and 5. tail laterally compressed with a distinct dorsal keel for almost the entire length of the tail, and the indistinct cross-stripes compressed only in the proximal half, the remainder of the tail uniformly yellowish (Fig. 6; Böhme, 2015). Examination of the skulls of the two species *Varanus griseus* and *Varanus nesterovi* by CT-scan and extraction of skull bones (the characteristics of the skull bones in






**Figure 5.** Photograph of a juvenile Bengal monitor lizard (*Varanus bengalensis*) captured in Bahu Kalat (Gandu) Protected Area (Oraie et al., 2024).



**FIGURE 6.** *Varanus nesterovi* in its natural environment at Mortka, Darbandikhan, Iraq (Böhme et al., 2023).

some lizards are a distinguishing feature between species) is true to some extent for the mentioned species; The similarity in the structure of the skull indicates a close relationship between the two species and their similar feeding method (Ghanbarinia et al., 2019). The documented localities of *Varanus nesterovi* form a distribution range that appears to be wedged between the respective ranges of *V. g. griseus* and *V. g. caspius*. As far as known, it is confined to the western and southwestern margin of the Zagros Mountain range on both sides of the Iraqi-Iranian border, extending down to the area of Shiraz, Fars Province, Iran. It seems that *V. nesterovi* prefers more elevated habitats, having been recorded from 600 to 1100 m (Fig. 5; Böhme, 2015). *V. nesterovi* is an endemic inhabitant of the southwestern Zagros Mountain range (Fig. 7). Its distribution area in submontane to montane steppe habitats is wedged between those of its two much more widespread relatives: *V. griseus* in the west and *V. caspius* in the east (Böhme et al., 2023).



**FIGURE 7.** *Varanus* distribution map in Iranian plateau. *V. nesterovi* (blue square), *V. griseus* (red circle), *V. bengalensis* (green circle) and *V. caspius* (red triangle). The presence points shown in this map are approximate. (The map was taken from the articles, Böhme et al., 2023; Oraie et al., 2024; Böhme, 2015; Fathnia et al., 2009; Shafiei Bafti et al., 2021; Yousefi et al., 2023; Nasrabadi et al., 2016; Ebrahimi Pour et al., 2016).

## CONCLUSION

The desert monitor lizard is the largest lizard in Iran, occurring across a wide area of the country. Consequently, maintaining suitable habitat for this species seems essential. Large areas in the northeast, south, and southwest of Iran are climatically favorable for the desert monitor lizard (Kafash, 2016). On the Iranian Plateau, four monitor species from two subgenera are present: *Psammosaurus* (including *Varanus griseus*, *Varanus caspius*, and *Varanus nesterovi*) and *Empagusia* (which includes *Varanus bengalensis*). *Varanus caspius* is an endangered species, and its distribution range has drastically changed, a trend expected to continue. Alarmingly, some protected areas have already become unsuitable habitats for this species (Shadloo et al., 2021). Furthermore, *V. bengalensis* has also been categorized as an endangered species (Oraie et al., 2024).

Given the similarity of the external and cranial features of *Varanus griseus* and *Varanus nesterovi*, as well as the proximity and, in some places, overlapping of their distribution areas, molecular studies are necessary to confirm their species separation. In recent years, we have witnessed the destruction of many animal habitats, including those of lizards, primarily due to human activities and climate fluctuations. Therefore, it seems necessary to examine the population dynamics, habitat status, and effects of climate fluctuations on other species. Further studies are also needed regarding their physiology, interspecific phylogeny, and other relevant aspects.

## ACKNOWLEDGMENTS

The authors thank Hossein Nabizadeh, and Morad Besharati for their assistance in this study.



## LITERATURE CITED

- Anderson, J., 1871. On some Indian reptiles. *Proceedings of the Zoological Society of London* 1871: 149–211
- Anderson, S. C., 1963. Amphibians and reptiles from Iran. *Proceedings of the California Academy of Sciences*, San Francisco State College.
- Anderson, S. C., 1999. The lizards of Iran. *Contributions to Herpetology* Volume 15 (vol. i–vii, pp. 1–442). Society for the Study of Amphibians and Reptiles. [Review in *Copeia*.2000;(4):1144.]
- Ast, J. C., 2001. Mitochondrial DNA evidence and evolution in Varanoidea (Squamata). *Cladistics*, 17(3), 211–226. <https://doi.org/10.1006/clad.2001.0169>
- Augé, M., 2005. Evolution des lézards du Paléogène en Europe. *Mémoires du Muséum National d'histoire Naturelle*, Paris, 192, 1–369.
- Arida E, Bo'hme W., 2010. The origin of *Varanus*: when fossils, morphology, and molecules alone are never enough. *Biawak*.; 4: 117–124
- Augé, M. L., Folie, A., Smith, R., Phélizon, A., Gigase, P., & Smith, T., 2022. Revision of the oldest varanid, *Saniwa orsmaelensis* Dollo, 1923, from the early-Eocene of northwest Europe. *Comptes Rendus Palevol*, 21, 511–529. <https://doi.org/10.5852/cr-palevol2022v21a25>
- Akman, B., Yıldız, M. Z., Aydoğdu, M., Bozkurt, M. A., & Üçeş, F., 2023. Species Conservation Action Plan for *Varanus griseus* (Daudin, 1803) Distributed in Southeastern Anatolia. *Forestist*, 73(2): 131–137. <https://doi.org/10.5152/forestist.2022.22027>
- Blanford, W. T., 1881. On a Collection of Persian Reptiles recently added to the British Museum. In *Proceedings of the Zoological Society of London* (Vol. 49, No. 3, pp. 671–682). Oxford, UK: Blackwell Publishing Ltd.
- Boettger, O., 1885. Liste der von Hrn. Dr. med. W. Kobelt in Algerien und Tunisien gesammelten Kriechtiere. *Reiseerinnerungen aus Algerien und Tunis. Berichte der Senckenbergischen Naturforschungs Gesellschaft*, 1885, 452–475.
- Boulenger, G.A., 1885. Catalogue of the lizards in the British Museum (Natural History). Vol. 2, Second edition. Printed by Taylor And Francis. London 1885.
- Boulenger, G. A., 1887. A list of the reptiles and batrachians obtained near Muscat, Arabia, and presented to the British Museum by Surgeon-Major ASG Jayakar. *Journal of Natural History*, 20(120), 407–408.
- Bohme, W., & Ziegler, T., 1997. A taxonomic review of the *Varanus* (Polydaedalus) niloticus (Linnaeus, 1766) species complex. *Herpetological Journal*, 7(4), 155–162.
- Bayless, M. K., 2002. Monitor lizards: A Pan-African check-list of their zoogeography (Sauria: Varanidae: P Polydaedalus). *Journal of Biogeography*, 29(12), 1643–1701. <http://doi.org/10.1046/j.1365-2699.2002.00779.x>
- Böhme, W., 2003. Checklist of the living monitor lizards of the world (family Varanidae). *Zoologische Verhandelingen*, 341, 4–43.

Böhme, W. & Ziegler, T., 2007. Notes on the Distribution, Diet, Hemipenis Morphology and Systematics of *Varanus spinulosus* Mertens, 1941. In: Horn, H.-G., Böhme, W. & Krebs, U. (Eds.), *Advances in Monitor Research III*, Mertensiella 16, pp. 100–108.

Baig, K. J., Masroor, R., & Arshad, M., 2008. Biodiversity and ecology of the herpetofauna of Cholistan Desert, Pakistan. *Russian Journal of Herpetology*, 15(3), 193–205

Buckitsch, Y., Bohme, W., Koch, A., 2016. Scale Morphology and Micro-Structure of Monitor Lizards (Squamata: Varanidae: *Varanus* spp.) and their Allies: Implications for Systematics, Ecology, and Conservation. *Zootaxa* 4153(1), 001–192. <http://doi.org/10.11646/zootaxa.4153.1.1>

Bauer, A. M., DeBoer, J. C., & Taylor, D. J., 2017. Atlas of the reptiles of Libya. *Proceedings of the California Academy of Sciences*, 64(8), 155–318. <http://doi.org/10.5281/zenodo.13155907>

Bar, A., Haimovitch, G., & Meiri, S., 2021. Field guide to the amphibians and reptiles of Israel. Frankfurt am Main, Allemagne. Frankfurt Contributions to Natural History, volume 87. 511 pages. <http://doi.org/10.48716/bullshf.182-3>

Brennan I. G., Lemmon A. R., Moriarty Lemmon E., Portik D. M., Weijola V., Welton L., Donnellan S. C., and Keogh J. S., 2021, Phylogenomics of monitor lizards and the role of competition in dictating body size disparity, *Systematic Biology*, 70(1), 120–132. <https://doi.org/10.1093/sysbio/syaa046>

Böhme, W., Ahmed, S. H., Al-Sheikhly, O. F., Ararat, K., Auer, M., Khudur, F. A., & Langner, C., 2023. Desert Monitor Lizards (Squamata: Varanidae: *Varanus*: *Psammosaurus*) from the Middle East: Further Records of Nesterov's Desert Monitor, *Varanus* (P.) *nesterovi* Böhme, Ehrlich, Milto, Orlov et Scholz, 2015, from Iraq, and Adjacent Localities of *Varanus* (P.) *g. griseus* (Daudin, 1803) and *Varanus* (P.) *g. caspius* (Eichwald, 1831), with Comments on Biogeography and Taxonomy. *Russian Journal of Herpetology*, 30(6), 518–528. <http://doi.org/10.30906/1026-2296-2023-30-6-518-52>

Calori, L., 1857. Sullo scheletro del Monitor terrestris Aegypti Cuv. (*Varanus arenarius* Duméril e Bibron). *Memorie della Reale Accademia delle Scienze dell'Istituto di Bologna*, 8, 161–202.

Conrad J.L., 2008. Phylogeny and systematics of Squamata (Reptilia) based on morphology. *Bulletin of the American Museum of Natural History* 310(1):1–182. <http://doi.org/10.1206/310.1>

Collar, D.C., Schulte II, J.A. & Losos, J.B., 2011. Evolution of extreme body size disparity in monitor lizards (*Varanus*). *Evolution*, 65, 2664–2680. <https://doi.org/10.1111/j.1558-5646.2011.01335.x>

Conrad J.L., Ast J.C., Montanari S., Norell M.A., 2011. A combined evidence phylogenetic analysis of Anguimorpha (Reptilia: Squamata). *Cladistics* 27:230–277. <https://doi.org/10.1111/j.1096-0031.2010.00330.x>

Chirikova M., Zima Yu. A., Pestov M. V., Gubin B. M., and Kovalenko A. V., 2019. Investigations on distribution, occurrence and threats of *Varanus griseus caspius* (Eichwald, 1831) at the north-eastern periphery of its geographical range (Kazakhstan), *Russian Journal of Herpetology*, 26(4), 185–200. <http://doi.org/10.30906/1026-2296-2019-26-4-185-200>

Daudin, F. M., 1802. Histoire Naturelle, générale et particulière des reptiles, ouvrage faisant suite, à l'histoire naturelle, générale et particulière composée par LECLERC DE BUFFON, et redigée par C. S. SONNINI, vol. 3. F. Dufart, Paris, 452 pp.

Daudin, F. M., 1803. Histoire Naturelle. Générale et Particulière des Reptiles, Vol. 8. F. Dufart, Paris, 439 pp.

Duméril, A.M. C., & G. Bibron., 1836. *Erpetologie Générale ou Histoire Naturelle Complete des Reptiles*. Vol. 3. Libr. Encyclopédique Roret, Paris, 528 pp

de Fejérváry, G. J., 1935. Further contributions to a monograph of the Megalanidae and Fossil Varanidae, with notes on recent *varanians*. *Annales Historico-Naturales Musei Nationalis Hungarici, Pars Zoologica*, 29, 1–130

Damadi, E., Karamiani, R., Rastegar-Pouyani, N., & Gholamifard, A., 2017. A checklist of lizards from southeastern part of the Sistan and Baluchestan Province in southeastern Iran. *Iranian Journal of Animal Biosystematics*, 13(1), 67–76. <https://doi.org/10.22067/ijab.v13i1.50398>

Eichwald, E., 1831. *Zoologia specialis, quam expositis animalibus tum vivis, tum fossilibus potissimum rossiae in universum, et poloniae in specie, in usum lectionum publicarum in Universitate Caesarea Vilnensi*.

Escoriza, D., Ferrer, J., Amat, F., & Tarragó, A., 2011. Nova cita I aspects ecològics del varà del desert *Varanus griseus* (Sauria: Varanidae) al Sahara Occidental. *Bulletin Society Catalana d'Herpetology*, 19, 70–74.

Ehrlich K., 2013. Morphologische Variation, Systematik und Biogeographie des Wüstenwarans, *Varanus griseus* (Daudin, 1803). MSc Thesis, University of Bonn.

Ebrahimi Pour, F., Rastegar-Pouyani, E., & Ghorbani, B., 2016. A preliminary study of the Reptile's fauna in northwestern Yazd province, Iran. *Russian Journal of Herpetology*, 23(4), 243–248.

Esquerre D., Donnellan S., Brennan I.G., Lemmon A.R., Lemmon E.M., Zaher H., Grazziotin F., Keogh J.S., 2020. Phylogenomics, biogeography, and morphometrics reveal rapid phenotypic evolution in pythons after crossing Wallace's line. *Systematic Biology*, Vol 69, Pages 1039–1051. <https://doi.org/10.1093/sysbio/syaa024>

Fathnia, B., Rastegar, P. N., Sampour, M., Bahrami, A. M., & Jaafari, G., 2009. The lizard fauna of Ilam province, Southwestern Iran. *Iranian Journal of Animal Biosystematics*, 5: 65–79. <https://doi.org/10.22067/ijab.v5i2.3340>

Geoffroy Saint-Hilaire, E., 1827. Description des Reptiles. In: Savigny, M. J. C. L. de, Description de l'Égypte ou recueil des observations et des recherches Quintana Roo ont été faites en Égypte pendant l'expédition de l'Armée Française, publié par les ordres de sa Majesté l'Empereur Napoléon Le Grand. Histoire Naturelle. Paris, L'imprimerie impériale. pp. 121–160.

Gray, J. E., 1845. Catalogue of the Specimens of Lizards in the Collection of the British Museum. order of the Trustees.

Gardner, A. S., 2013. The Amphibians and Reptiles of Oman and the UAE. Edition Chimaira.

Georgalis, G. L., 2017. Necrosaurus or *Palaeoivanus*? Appropriate nomenclature and taxonomic content of an enigmatic fossil lizard clade (Squamata). *Annales de Paléontologie*, 103, 293–303. <https://doi.org/10.1016/j.annpal.2017.10.001>

Ghanbarinia, N., Rastegar-Pouyani, N., & Karamiani, R., 2019. A Study of Skull Structure in the Desert Monitor, *Varanus griseus* (Daudin, 1803) (Sauria: Varanidae). *Taxonomy and Biosystematics*, 11(40), 59–74. <https://doi.org/10.22108/tbj.2020.123740.1117>

- Georgalis, G. L., Čerňanský, A., & Klembara, J., 2021. Osteological atlas of new lizards from the Phosphorites du Quercy (France), based on historical, forgotten, fossil material. *Geodiversitas*, 43, 219–293. <http://doi.org/10.5252/geodiversitas2021v43a9>
- Georgalis, G. L., Menecart, B., & Smith, K. T., 2023. First fossil record of *Varanus* (Reptilia, Squamata) from Switzerland and the earliest occurrences of the genus in Europe. *Swiss Journal of Geosciences*, 116(1), 9. <http://doi.org/10.1186/s00015-023-00440-5>
- Hofstetter, R., 1969. Présence de Varanidae (Reptilia, Sauria) dans le Miocène de Catalogne. Considérations sur l'histoire de la famille. *Bulletin du Muséum National d'histoire Naturelle*, 40, 1051–1064.
- Heidari, N., & Kami, H. G., 2009. Lizards of the Gando Protected area in Sistan and Baluchestan province, southeastern Iran. *Iranian Journal of Animal Biosystematics*, 5: 57–64 <https://doi.org/10.22067/ijab.v5i2.3339>
- Holmes, R., Murray, A., Attia, Y., 2010. Oldest known *Varanus* (Squamata: Varanidae) from the upper eocene and lower oligocene of Egypt: support for an African origin of the genus. *Palaeontology*, Vol. 53, Part 5, 2010, pp. 1099–1110. <https://doi.org/10.1111/j.1475-4983.2010.00994.x>
- Ivanov, M., Ruta, M., Klembara, G., Böhme, M., 2017. A new species of *Varanus* (Anguimorpha: Varanidae) from the early Miocene of the Czech Republic, and its relationships and palaeoecology. *Journal of Systematic Palaeontology*. <http://doi.org/10.1080/14772019.2017.1355338>
- Jablonski, D., Bommel, S., & Masroor, R., 2021. Notes on the herpetofauna of Kunduz Province, northeastern Afghanistan, with the first record of *Varanus griseus caspius* (Eichwald, 1831). *Herpetology Notes*, 14, 63–66.
- Jaman, M. F., Ahmed, S., Hossain, S., & Rabbe, M. F., 2024. Activity budget of Bengal Monitor *Varanus bengalensis*: Effect of daytime, season, age and temperature. *Heliyon*, 10(7). <https://doi.org/10.1016/j.heliyon.2024.e28504>
- Keogh J.S., 1998. Molecular phylogeny of elapid snakes and a consideration of their biogeographic history. *Biological Journal of the Linnean Society*, 63:177–203. <https://doi.org/10.1006/bjil.1997.0178>
- Khan, M. S., 2003. Checklist and key to the lizards of Pakistan. *Pakistan Journal of Zoology*.(Suppl. Ser.), 5, 1–25.
- Khan, M. S., 2004. Annotated checklist of amphibians and reptiles of Pakistan. *Asiatic Herpetological Research*, 10, 191–201.
- Kabir, S. M. H., Ahmad, M., Ahmed, A. T. A., Rahman, A. K. A., Ahmed, Z. U., Begum, Z. N. T., ... & Khandker, M., 2009. Encyclopedia of flora and fauna of Bangladesh. Vol-25. *Amphibians and Reptiles*, 1–204.
- Koch, A., Auliya, M. & Ziegler, T., 2010. Updated checklist of the living monitor lizards of the world (Squamata: Varanidae). *Bonn Zoological Bulletin*, 57(2), 127–136
- Koch, A., Ziegler, T., Boehme, W., Arida, E., & Auliya, M., 2013. Pressing problems: Distribution, threats, and conservation status of the monitor lizards (Varanidae: *Varanus* spp.) of Southeast Asia and the Indo-Australian Archipelago. *Herpetological Conservation and Biology*, 8(3), 1–62.
- Kafash, A., 2016. Investigation of the most important effective climate variable on the presence of *Varanus griseus* species and its limiting climatic variables. *Zist Sepehr Student Magazine*, 10(2), 30–35.



- Levshakova, I. J., 1986. A new *Varanus* from lower Pliocene of Tajikistan. *Trudy Zoologicheskogo Instituta Akademii Nauk SSSR*, 157, 101–116.
- Merrem, B., 1820. Versuch eines systems der Amphibien. Krieger. Marburg, J.C. Kreiger, 1820, German. <https://doi.org/10.5962/bhl.title.5037>
- Mertens, R., 1942. Die Familie der Warane (Varanidae). Erster bis dritter Teil Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft, 462, 465, 466, 1–391.
- Mertens R., 1954. Über die Rassen des Wüstenwarans (*Varanus griseus*). *Senckenbergiana Biologica*, 35, 353–357.
- Mertens, R., & Wermuth, H., 1960. Die Amphibien und Reptilien Europas:(Dritte Liste, nach dem Stand vom 1. Januar 1960). Kramer.
- Mertens R., 1973. Zur Kenntnis des iranischen Wüstenwarans, *Varanus griseus*. *Bonner Zoologische Beiträge* 24, 233 – 236.
- Molnar, R. E., 2004. The long and honorable history of Monitors and their kin. In R. E. Pianka, D. King, & R. A. King (Eds.), *Varanoid Lizards of the World* (pp. 10–67). Indiana University Press.
- MacCoy M., 2015. A field guide to the reptiles of the Solomon Islands. Michael McCoy, Kuranda.
- Malakhov, D. V., & Chirikova, M. A., 2018. Species distribution model of *Varanus griseus caspius* (Eichwald, 1831) in Central Asia: An insight to the species' biology. *Russian Journal of Herpetology*, 25(3), 195–206. <http://doi.org/10.30906/1026-2296-2019-25-3-195-206>
- Nilson, G., & Andren, C. L. A. E. S., 1981. Die Herpetofauna des Kavir-Schutzgebietes, Kavir-Wüste, Iran. *Salamandra*, 17(3–4), 130–146.
- Nasrabadi, R., Rastegar Pouyani, E., Hosseinian Yousefkhani, S. S., & Khani, A., 2016. A checklist of herpetofauna from Sabzevar, Northeastern Iran. *Iranian Journal of Animal Biosystematics*, 12(2), 255–259. <https://doi.org/10.22067/ijab.v12i2.37650>
- Oraie, H., Mobaraki, A., & Abtin, E., 2024. Long-range relatives: on the status of the Bengal monitor, *Varanus bengalensis* (Squamata: Varanidae) from southeastern Iran. *Iranian Journal of Animal Biosystematics*, 20(1), 45–50. <https://doi.org/10.22067/ijab.2024.88144.1072>
- Pianka E.R., 1995. Evolution of body size: varanid lizards as a model system. *The American Naturalist*, 146:398–414.
- Pyron, R. A., Burbrink, F. T., & Wiens, J. J., 2013. A phylogeny and revised classification of Squamata, including 4161 species of lizards and snakes. *BMC Evolutionary Biology*, 13(1), 1–54. <https://doi.org/10.1186/1471-2148-13-93>
- Pyron R.A., 2017. Novel approaches for phylogenetic inference from morphological data and total-evidence dating in squamate reptiles (lizards, snakes, and amphisbaenians). *Systematic Biology*, 66:38–56. <https://doi.org/10.1093/sysbio/syw068>
- Rabl-Rückhard, H., 1881 Über das Vorkommen eines Fornixrudiments bei Reptilien. (No Title).

Rastegar-Pouyani, N., Johari, M., Rastegar-Pouyani, E., 2007. Field Guide to the Reptiles of Iran. Volume1: Lizards. Second edition. Iran, Razi University Publishing. 296 p. (In Farsi).

Reynolds G.R., Niemiller M.L., Revell L.J., 2014. Toward a Tree-of-Life for the boas and pythons: multilocus species-level phylogeny with unprecedented taxon sampling. *Molecular Phylogenetics and Evolution* 71:201–13. <http://doi.org/10.1016/j.ympev.2013.11.011>

Schlegel, H., 1839. Abbildungen neuer oder unvollständig bekannter Amphibien, nach der Natur oder dem Leben entworfen und mit einem erläuternden Texte begleitet. Arne and Comp.

Sixta, V., 1900. Vergleichend-osteologische Untersuchungen über den Bau des Schädels von Monotremen und Reptilien. *Zoologischer Anzeiger*, 23, 213–229.

Sindaco, R., Venchi, A., Carpaneto, G. M., & Bologna, M. A., 2000. The Reptiles of Anatolia: A Checklist and Zoogeographical analysis. *Biogeographia. The Journal of Integrative Biogeography*, 21, 441–554. <http://doi.org/10.21426/B6110017>

Stanner, M. 2004. *Varanus griseus*. In: E.R. Pianka and D. King (eds), *Varanoid Lizards of the World*, pp.104–132. Indiana University Press, Bloomington, Indiana.

ŠMÍD, J., Moravec, J., Kodym, P., Kratochvíl, L., Yousefkhani, S. S. H., & Frynta, D., 2014. Annotated checklist and distribution of the lizards of Iran. *Zootaxa*, 3855(1), 1–97. <http://dx.doi.org/10.11646/zootaxa.3855.1.1>

Safaei-Mahroo, B., Ghaffari, H., Fahimi, H., Broomand, S., Yazdanian, M., Najafi-Majd, E., ... & Kazemi, S. M., 2015. The herpetofauna of Iran: checklist of taxonomy, distribution and conservation status. *Asian Herpetological Research*, 6(4), 257–290. <http://doi.org/10.16373/j.cnki.ahr.140062>

Shadloo, S., Mahmoodi, S., Hosseinzadeh, M. S., & Kazemi, S. M., 2021. Prediction of habitat suitability for the desert monitor (*Varanus griseus caspius*) under the influence of future climate change. *Journal of Arid Environments*, 186, 104416. <http://doi.org/10.1016/j.jaridenv.2020.104416>

Shafiei Bafti, S., Rouhani, S., Sehhatiasabet, M. E., Moradi, N., & Mansour Mirtadzadini, S., 2021. A Study of Reptiles Fauna of the Saadi Protected Area in Kerman Province. *Taxonomy and Biosystematics*, 13(48), 93–108. <https://doi.org/10.22108/tbj.2022.130988.1183>

Theobald, W., 1868. Catalogue of the Reptiles of British Birma, embracing the Provinces of Pegu, Martaban, and Tenasserim; with descriptions of new or little-known species. *Zoological Journal of the Linnean Society*, 10(41), 4–67. <https://doi.org/10.1111/j.1096-3642.1868.tb02007.x>

Tuck, R. G., 1971. Amphibians and reptiles from Iran in the United State National Museum Collection. *Bulletin of the Maryland Herpetological Society*, 7(3), 48–36.

Tsellarius, A. Y., Tsellarius, E. Y., & Men'shikov, Y. G., 1997. Notes on the diet and foraging of *Varanus griseus*. *Russian Journal of Herpetology*, 4(2), 170–181. <https://doi.org/10.30906/1026-2296-1997-4-2-170-181>

Tsellarius, A. Y., & Tsellarius, E. Y., 1997. Behavior of *Varanus griseus* during encounters with conspecifics. *Asiatic Herpetological Research*, 7, 108–130.

Uetz, P., Freed, P., Aguilar, R., Reyes, F., Kudera, J. & Hošek, J. (eds.) (2025) The Reptile Database, <http://www.reptile-database.org>, accessed [10 April 2025].

Vitt, L.J., Caldwell, J.P., 2013. Herpetology: an introductory biology of amphibians and reptiles. Academic press. <http://doi.org/10.11606/issn.2316-9079.v12i2p147-149>

Voris, H., 2000. Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time durations. *Journal of Biogeography*, 27, 1153–1167. <https://doi.org/10.1046/j.1365-2699.2000.00489.x>

Villa, A., Abella, J., Alba, D. M., Almécija, S., Bolet, A., Koufos, G. D., Knoll, F., Luján, À. H., Morales, J., Robles, J. M., Sánchez, I. M., & Delfino, M., 2018. Revision of *Varanus marathonensis* (Squamata, Varanidae) based on historical and new material: Morphology, systematics, and paleobiogeography of the European monitor lizards. *PLoS ONE*, 13, e0207719. <https://doi.org/10.1371/journal.pone.0207719>

Wilson, D. S., 1975. The adequacy of body size as a niche difference. *The American Naturalist*, 109(970), 769–784.

Weijola, V., 2017. Phylogeny and Taxonomy of the Pacific Monitor Lizards (Squamata: *Varanus*: Euprediosaurus). University of Turku, Turku.

Yousefi, M., Nezami, B., Adibi, M. A., Geray, A., & Ataei, F., 2023. Reptile richness in Touran Biosphere Reserve, Semnan province. *Experimental animal Biology*, 12(1), 77–85. <https://doi.org/10.30473/eab.2023.59681.1838>

Ziegler, T., Schmitz, A., Koch, A. & Böhme, W., 2007. A review of the subgenus Euprepiosaurus of *Varanus* (Squamata: Varanidae): morphological and molecular phylogeny, distribution and zoogeography, with an identification key for the members of the *V. indicus* and the *V. prasinus* species groups. *Zootaxa*, 1472, 1–28. <https://doi.org/10.11646/zootaxa.1472.1.1>