

RESEARCH ARTICLE

Open access

# Checklist of Collembola (Hexapoda) from Iran, Part II: Entomobryomorpha

Masoumeh Shayanmehr<sup>1,\*</sup>, Rafael Jordana<sup>2</sup>, Elham Yoosefi Lafooraki<sup>3</sup>, Eliyeh Yahyapour<sup>4</sup>, Mehdi Ghajar Sepanlou<sup>5</sup>

<sup>1</sup>Department of Plant Protection, Faculty of Crop Sciences, Sari University of Agricultural Sciences and Natural Resources (SANRU), Mazandaran province, Iran. [m.shayanmehr@sanru.ac.ir](mailto:m.shayanmehr@sanru.ac.ir)

<sup>2</sup>University of Navarra, Department of Environmental Biology, Pamplona, Spain. [rjordana@unav.es](mailto:rjordana@unav.es)

<sup>3</sup>Department of Plant Protection, Faculty of Crop Sciences, Sari University of Agricultural Sciences and Natural Resources (SANRU), Mazandaran province, Iran. [elham.yoosefi@sanru.ac.ir](mailto:elham.yoosefi@sanru.ac.ir)

<sup>4</sup>Islamic Azad University, Arak branch, Agricultural Faculty, Entomology Department, P.O. Box 38135/567, Arak, Iran. [eyahyapur@yahoo.com](mailto:eyahyapur@yahoo.com)

<sup>5</sup>Department of Soil Science, Faculty of Crop Sciences, Sari University of Agricultural Sciences and Natural Resources (SANRU), Mazandaran province, Iran. [sepanlou@gmail.com](mailto:sepanlou@gmail.com)

## Abstract

The order Entomobryomorpha is one of the four important orders of the Collembola (Springtails). These arthropods are distinguished from the rest of the Collembola by having the third abdominal segment larger than the fourth segment and having scales in some species. In this study, the list of Entomobryomorpha species (Collembola) in Iran has been prepared based on scientific sources related to Collembola reports between 2020 and 2025 and the materials examined in the present study. This checklist includes 139 species in 41 genera belonging to six families of the order Entomobryomorpha along with the materials examined and distribution in the world and Iran. Three species, including one species from the family Entomobryidae, *Lepidocyrtus lignorum* (Fabricius, 1775), one species from the family Tomoceridae, *Tomocerus asiaticus* Martynova 1969, and the species *Desoria kaszabi* Dunger, 1982 from the family Isotomidae, are new to the fauna of Iran. A list of Entomobryomorpha species, descriptions of new records, along with some images and a key to the Iranian species of the family Tomoceridae are provided.

**Keywords:** Mazandaran, Mesofauna, soil-living animals, taxonomy.

**Received:**  
09 April 2025

**Accepted:**  
02 July 2025

## INTRODUCTION

Order Entomobryomorpha Börner, 1913 (Collembola), the elongate-bodied springtails, are ubiquitous, species-rich, and diversified hexapods (Yu et al., 2022). They have long and slim bodies unlike Symphyleona and Neelipleona orders which are known as globular springtails and a reduced prothorax unlike Poduromorpha order which have a well-developed prothorax (Hopkin, 1997). The members of main groups of Entomobryomorpha are cosmopolitan so that occur in any terrestrial niches (Yu et al., 2022).

The study of Collembola fauna in Iran has been increasing over the last few years so that several species have been described or recorded from the country. As a result, the endemism among Collembola species of Iran is increasing. The list of Iranian springtails published by Cox (1982) includes 31 species in 13 genera of Entomobryomorpha. In the checklist of Collembola species of Iran by Shayanmehr et al. (2013), 56 species in 23 genera of Entomobryomorpha were reported, and this number were increased to 105 species in 34 genera in the next list (Shayanmehr et al., 2020b). Present work deals with the list of

\*Corresponding Author: ✉ : [m.shayanmehr@sanru.ac.ir](mailto:m.shayanmehr@sanru.ac.ir)



Entomobryomorpha species based on published literature between 2020 and 2024 and the materials of present study.

## MATERIAL AND METHODS

This checklist is based on the literature published between 2020 and 2025 and collected materials by authors. Leaf litter, soil, and dead wood were collected from various localities in Mazandaran province, North of Iran, specimens were extracted using Tullgren funnels and preserved in 75–80% alcohol. To prepare microscopic slides samples were cleared in 10% KOH and mounted on Hoyer's medium. The identification of specimens was carried out using valid identification keys such as Potapov (2001), Fjellberg (2007), Jordana (2012), and Winkler et al. (2020). The examined materials are deposited in Entomology lab of Sari University of Agricultural Sciences and Natural Resources, Iran.

## RESULTS

The checklist of Entomobryomorpha order of Iran is provided here alphabetically. In total, this order currently comprises 139 species, 41 genera, and six springtails families recorded in Iran. Three species are newly recorded for Iran which are denoted by an asterisk (\*).

**Order Entomobryomorpha Börner, 1913**  
**Family Entomobryidae Schäffer, 1896**  
**Subfamily Entomobryinae Schäffer, 1896**  
**Genus *Calx* Christiansen, 1958**

***Calx kailashi* Mandal, 2018**

**General distribution:** India and Iran (Mandal, 2018; Ahmadi et al., 2023).

**Distribution in Iran:** Kurdistan (Marivan-Pirsafa) province (Ahmadi et al., 2023).

**Genus *Coecobrya* Yosii, 1956**

***Coecobrya caeca* (Schött, 1896)**

**General distribution:** Australia, North America (Greenslade, 1992).

**Distribution in Iran:** Kerman (Mahan) province as *Sinella* cf. *coeca* (Schött, 1896) (Abdolalizadeh, 2018).

***Coecobrya tenebricosa* (Folsom, 1902)**

**General distribution:** Cosmopolitan (Jordana, 2012).

**Distribution in Iran:** Isfahan and Mazandaran (Babol) provinces as *Sinella tenebricosa* Folsom, 1902 (Nematollahi et al., 2009; Alijani-Ardeshtir et al., 2017).

**Genus *Drepanura* Schött, 1891**

***Drepanura kirgisica* Martynova, 1971**

**General distribution:** Central and North Asia (Abdolalizadeh et al., 2019).

**Distribution in Iran:** Kerman province (Abdolalizadeh, 2018; Abdolalizadeh et al., 2019).

***Drepanura pallens* Rusek, 1981**

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** Fars province, synanthropic species (Semati et al., 2023).

**Examined material:** Mazandaran province, Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 2.

***Drepanura tuxeni*** Nosek, 1964**General distribution:** Palaearctic, Tajikistan (Rabieh et al., 2023).**Distribution in Iran:** South Khorasan (Birjand) province (Rabieh et al., 2023)**Genus *Entomobrya*** Rondani, 1861***Entomobrya albocincta*** (Templeton, 1836) in Templeton & Westwood 1836**General distribution:** Europe, Mediterranean (Bellinger et al., 2024).**Distribution in Iran:** Kohgiluyeh and Boyer-Ahmad (Cheram) province (Falahati Hosseiniabad et al., 2011).***Entomobrya atrocincta*** Schött, 1896**General distribution:** Europe, Mediterranean (Bellinger et al., 2024).**Distribution in Iran:** Golestan (Gorgan), Kermanshah (Harsin), Mazandaran (Behshahr, Qaemshahr, Sari, Savadkooh) and Tehran provinces (Yahyapour et al., 2011; Yoosefi Lafooraki, 2014; Kahrarian, 2015, 2019; Darvish-Motevalli, 2016; Hosseini et al., 2016a; Qazi & Shayanmehr, 2016; Yahyapour et al., 2020; Ghasemi Charati et al., 2022; Vahedi Moghadam et al., 2022).***Entomobrya corticalis*** (Nicolet, 1842)**General distribution:** Palaearctic (Bellinger et al., 2024).**Distribution in Iran:** Golestan (Golestan National Park), Guilan, Mazandaran (Sari), and North Khorasan (Darkesh-Jozak forest, Zoeram village) provinces (Cox, 1982; Khanahmadi, 2018; Yahyapour et al., 2018, 2019; Mehrafrooz Mayvan et al., 2022).**Examined material:** Mazandaran province, Behshahr, Zenget village, Hezarjirib forest, 36°25'15" N, 53°34'16" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 3; Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 2.***Entomobrya handschini*** Stach, 1922**General distribution:** Europe, Poland to Iran to the south east of Black Sea (Baquero et al., 2008).**Distribution in Iran:** Kermanshah (Harsin, Kermanshah, Sarab-e-Nilufar) and Tehran provinces (Moravvej, 2003; Qazi & Shayanmehr, 2014; Kahrarian et al., 2014; Kahrarian, 2015, 2019).***Entomobrya indica*** (Baijal, 1955)**General distribution:** Tibet, India (Jordana, 2012, Yadav et al., 2018).**Distribution in Iran:** Kerman (Sirch village) province (Abdolizadeh, 2018; Abdolizadeh et al., 2019).***Entomobrya lanuginosa*** (Nicolet, 1842)**General distribution:** Central and Northern Europe (Jordana, 2012).**Distribution in Iran:** Central, East Azarbaijan, Guilan, Mazandaran, and North Khorasan (Darkesh-Jozak forest) provinces (Cox, 1982; Mehrafrooz Mayvan et al., 2022).***Entomobrya lindbergi*** Stach, 1960**General distribution:** Palaearctic (Bellinger et al., 2024).**Distribution in Iran:** Guilan (Rasht), Kermanshah (Derkah, Harsin, Sarab-e Nilufar, Sahneh), Mazandaran (Sari), and Tehran provinces (Daghighi et al., 2013; Moravvej, 2003; Qazi & Shayanmehr, 2014; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Kahrarian, 2015, 2019; Kahrarian et al., 2014).**Examined material:** Mazandaran province, Behshahr, Zenget village, Hezarjirib forest, 36°25'15" N, 53°34'16" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 4; Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 3; Neka, Lemon

village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 2; Sari, Marzrud village, 36°14'20" N, 53°39'10" E, 46 m a.s.l., 18-May-2020, leg. E. Yahyapour, n= 81; Sari, Abmal forest, 36°42'52" N, 53°06'00" E, 10 m a.s.l., 24-May-2020, leg. E. Yahyapour, n= 35; Sari, Goharbaran seashore, soil under eucalyptus, 36°73'12" N, 53°11'14" E, 5 m a.s.l., 12-May-2020, leg. E. Yahyapour, n= 6; Sari, Dashtenaz forest, 36°42'10"N, 53°9'11" E, 11 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 6; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 6; Sari, rice field, 36°42'11"N, 53°10'12" E, 11 m a.s.l., 12-May-2020, leg. E. Yahyapour, n= 56.

***Entomobrya mesopotamica*** Rusek, 1981

**General distribution:** Iraq (Jordana & Baquero 2010).

**Distribution in Iran:** Kermanshah (Gahvareh, Sarabe-bid, Sorkh) and Markazi (Arak) provinces (Shoeibi et al., 2013; Kahrarian et al., 2014; Kahrarian, 2015).

***Entomobrya multifasciata*** (Tullberg, 1871)

**General distribution:** Holarctic (Babenko et al., 2019).

**Distribution in Iran:** Golestan (Gorgan, Kordkuy), Kurdistan (Marivan-Garan, Marivan-Pirsafa), and Mazandaran (Babol, Galugah, Juybar, Qaemshahr, Sari) provinces (Yahyapour et al., 2011; Yoosefi Lafooraki, 2014; Darvish-Motevalli, 2016; Hosseini et al., 2016; Alijani-Ardeshtir et al., 2017; Ghasemi Charati et al., 2022; Ahmadi et al., 2023).

***Entomobrya muscorum*** (Nicolet, 1842)

**General distribution:** Subcosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Kohgiluyeh and Boyer-Ahmad (Cheram) province (Falahati Hosseinabad et al., 2011).

***Entomobrya nicoleti*** (Lubbock, 1868)

**General distribution:** Palaearctic (Fjellberg, 2007).

**Distribution in Iran:** Golestan (Gorgan, Kordkuy) province (Hosseini et al., 2016).

***Entomobrya nigrocincta*** Denis, 1923

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** Kermanshah (Char Zabar, Harasam, Qal'eh) and Mazandaran (Neka, Sari, Savadkooh) provinces (Kahrarian, 2015; Kahrarian et al., 2014; Yahyapour et al., 2018; Bakhshi et al., 2022; Zamani Khormandichali et al., 2024).

**Examined material:** Mazandaran province, Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 2; Sari, Goharbaran seashore, soil under eucalyptus, 36°42'12"N 53°11'14"E, 5 m a.s.l., 12-May-2020, leg. E. Yahyapour, n= 4.

***Entomobrya nivalis*** (Linnaeus, 1758)

**General distribution:** Northern Holarctic (Babenko et al., 2019).

**Distribution in Iran:** Golestan (Gorgan, Kordkuy) province (Hosseini et al., 2016).

***Entomobrya numidica*** Baquero, Hamra-Kroua & Jordana, 2009

**General distribution:** Algeria (Jordana, 2012).

**Distribution in Iran:** Golestan (Gorgan) province (Hosseini et al., 2016).

***Entomobrya obscurella*** Brown, 1926

**General distribution:** Afghanistan, Palestine, Iraq (Jordana, 2012).

**Distribution in Iran:** Mazandaran (Sari, Savadkooh) province (Ghasemi Charati, 2017; Ghasemi Charati et al., 2022; Zamani Khormandichali et al., 2024).

*Entomobrya quinquelineata* Börner, 1901

**General distribution:** Europe (Jordana, 2012).

**Distribution in Iran:** Markazi (Arak) province (Shoeibi et al., 2013).

*Entomobrya schoetti* Stach, 1922

**General distribution:** Europe (Jordana, 2012).

**Distribution in Iran:** Guilan and Mazandaran (Neka) provinces (Kahrarian et al., 2014; Kahrarian, 2015; Yahyapour et al., 2018).

*Entomobrya unostrigata* Stach, 1930

**General distribution:** Mediterranean, Asia and possibly in USA (Jordana, 2012).

**Distribution in Iran:** Tehran province (Moravvej, 2003).

#### **Genus *Mesentotoma* Salmon, 1942**

*Mesentotoma dollfussi* (Denis, 1924)

**General distribution:** Europe, Algeria, Macaronesia (Jordana, 2012).

**Distribution in Iran:** Mazandaran (Qaemshahr) province (Darvish-Motevalli, 2016).

*Mesentotoma subdollfusi* Jacquemart, 1974

**General distribution:** Tunisia (Jordana, 2012).

**Distribution in Iran:** Golestan (Gorgan, Kordkouy), Guilan (Rasht), and Mazandaran (Qaemshahr, Savadkooh) provinces (Daghighi et al., 2013; Hosseini et al., 2016; Darvish-Motevalli, 2016; Zamani Khormandichali et al., 2024).

#### **Genus *Prodrepanura* Stach, 1963**

*Prodrepanura* sp.

**Distribution in Iran:** Kermanshah province (Kahrarian et al., 2012).

#### **Genus *Sinella* Brook, 1882**

*Sinella curviseta* Brook, 1882

**General distribution:** Probably Holarctic (Jordana, 2012).

**Distribution in Iran:** Kerman (Shahdad), Mazandaran (Babol, Qaemshahr, Savadkooh), and Tehran provinces (Moravvej, 2003; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Darvish-Motevalli, 2016; Alijani-Ardeshtir et al., 2017; Abdolalizadeh, 2018).

#### **Subfamily Lepidocyrtinae Wahlgren, 1906**

##### **Genus *Lepidocyrtus* Bourlet, 1839**

*Lepidocyrtus bicoloris* Mateos, 2012

**General distribution:** Spain (Mateos, 2012).

**Distribution in Iran:** Mazandaran (Neka) province (Bakhshi et al., 2022).

**Examined material:** Mazandaran province, Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 1; Neka, Sapelmiun village, 36°23'11" N, 53°34'18" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 10; Sari, Abmal forest, 36°42'52" N, 53°06'00" E, 10 m a.s.l., 24-May-2020, leg. E. Yahyapour, n= 1.

*Lepidocyrtus curvicolis* Bourlet, 1839

**General distribution:** Subcosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Kohgiluyeh and Boyer-Ahmad (Cheram) province (Falahati Hosseinabad et al., 2011).

*Lepidocyrtus cyaneus* Tullberg, 1871

**General distribution:** Cosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Central, East Azarbaijan, Golestan (Gorgan, Kordkuy), Guilan, Mazandaran, and Zanzan provinces (Cox, 1982; Hosseini et al., 2016).

**Examined material:** Mazandaran province, Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 4.

*Lepidocyrtus lanuginosus* (Gmelin, 1788)

**General distribution:** Subcosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Central, East Azarbaijan, Golestan (Gorgan), Guilan, and Mazandaran (Neka) provinces (Cox, 1982; Hosseini et al., 2016; Bakhshi et al., 2022).

**Examined material:** Mazandaran province, Neka, Sapelmiun village, 36°23'11" N, 53°34'18" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 1; Sari, Abmal forest, 36°42'52" N, 53°06'00" E, 10 m a.s.l., 24-May-2020, leg. E. Yahyapour, n= 2; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 1.

*Lepidocyrtus lignorum* (Fabricius, 1775)\* (Figures 1A, B)

**General distribution:** Subcosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** The species is reported for the first time from Iran.

**Material examined:** Mazandaran province, Behshahr, Zenget village, Hezarjirib forest, 36°25'15" N, 53°34'16" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 1.

*Lepidocyrtus paradoxus* Uzel, 1890

**General distribution:** Subcosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Kohgiluyeh and Boyer-Ahmad (Cheram) province (Falahati Hosseinabad et al., 2011).

*Lepidocyrtus ruber* Schött, 1902

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** Central, East Azarbaijan, Guilan, and Mazandaran provinces (Cox, 1982).

**Genus *Pseudosinella* Schäffer, 1897***Pseudosinella alba* (Packard, 1873)

**General distribution:** Cosmopolitan (Fjellberg, 2007).

**Distribution in Iran:** Guilan (Rasht) and Kurdistan (Marivan-Garan, Marivan-Pirsafa) provinces (Daghighi et al., 2013; Ahmadi et al., 2023).

**Examined material:** Mazandaran province, Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 1.

*Pseudosinella baghdadica* Rusek, 1981

**General distribution:** Iraq (Rusek, 1981).

**Distribution in Iran:** Kermanshah (Derkah, Gerdeh Now, Harsin, Quri Qal'eh, Sahneh, Satar, Sonqor, Tamark) and Khuzestan (Shush) provinces (Yoosefi Lafooraki et al., 2014; Kahrarian, 2015; Kahrarian et al., 2014).




FIGURE 1. A) Habitus of *Lepidocyrtus lignorum* (Magnification 40X); B) Labium chaetotaxy of *Lepidocyrtus lignorum*; R\* half in length than other setae (Magnification 100X).

***Pseudosinella cf. decipiens* Denis, 1924**

**General distribution:** Palaearctic (Fjellberg, 2007).

**Distribution in Iran:** Mazandaran (Sari) province (Shayanmehr et al., 2024).

***Pseudosinella duodecimpunctata* Denis, 1931**

**General distribution:** Europe (Bellinger et al., 2024).

**Distribution in Iran:** East Azarbaijan province (Cox, 1982).

***Pseudosinella immaculata* (Lie Pettersen, 1896)**

**General distribution:** Palaearctic (Fjellberg, 2007).

**Distribution in Iran:** Mazandaran (Sari) province (Shayanmehr et al., 2024).

***Pseudosinella imparipunctata* Gisin, 1953**

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** Guilan, Mazandaran, and Zanzan provinces (Cox, 1982).

***Pseudosinella octopunctata* Börner, 1901**

**General distribution:** Cosmopolitan (Fjellberg, 2007).

**Distribution in Iran:** Central, East Azarbaijan, Golestan (Gorgan, Kordkuy), Guilan, Ilam, Isfahan (Zarrinshahr), Kerman (Shahdad), Kermanshah (Eslamabad-e-gharb, Gahvareh, Guilan-e-Gharb, Harsin, Javanrud, Kangavar, Kermanshah, Paveh, Qasr-e-shirin, Sahneh, Sar-e-pol-e- Zahab, Sonqor, Sonqor, Tazeh abad), Kohgiluyeh and Boyer-Ahmad, Kurdistan (Marivan-Garan, Marivan-Pirsafa), Lorestan, Markazi, Mazandaran (Babol, Babolsar, Behshahr, Larijan, Qaemshahr, Sari, Savadkooh), North Khorasan (Darkesh-Jozak forest, Golil-Sarany rangeland, Oghaz village), Tehran, West Azarbaijan, and Zanzan provinces (Cox, 1982; Yahyapour et al., 2011; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Qazi & Shayanmehr, 2014; Kahrarian et al. 2014; Balvasi et al., 2015; Kahrarian, 2015; Darvish-Motevalli, 2016; Hosseini et al., 2016; Alijani-Ardeshtir et al., 2017; Abdolalizadeh, 2018; Ghasemi Charati, 2017; Moradi et al., 2018; Ghasemi Charati et al., 2022; Mehrafrooz Mayvan et al., 2022; Vahedi Moghadam et al., 2022; Ahmadi et al., 2023; Zamani Khormandichali et al., 2024).

**Examined material:** Mazandaran province, Behshahr, Zenget village, Hezarjirib forest, 36°25'15" N, 53°34'16" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 9; Neka, Lemon village, Hezarjirib

forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 4; Sari, Goharbaran seashore, soil under eucalyptus, 36°42'12"N 53°11'14"E, 5 m a.s.l., 12-May-2020, leg. E. Yahyapour, n=1.

***Pseudosinella sexoculata* Schött, 1902**

**General distribution:** Cosmopolitan (Fjellberg, 2007).

**Distribution in Iran:** Kermanshah and Kurdistan (Marivan-Garan, Marivan-Pirsafa) provinces (Ghahramaninezhad et al., 2013; Kahrarian, 2015; Ahmadi et al., 2023).

**Subfamily Seirinae Yosii, 1961**

**Genus *Seira* Lubbock, 1870**

***Seira atlantica* Negri, Pellicchia & Fanciulli, 2005**

**General distribution:** Morocco (Negri et al., 2005).

**Distribution in Iran:** North Khorasan (Varaghi village) and Zanzan provinces (Shayanmehr et al., 2020a,b; Mehrafrooz Mayvan et al., 2022).

***Seira domestica* (Nicolet, 1842)**

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** Golestan (Gorgan, Kordkouy), Guilan (Rasht), Kermanshah (Harsin), and Mazandaran (Sari, Savadkooh, Qaemshahr) provinces (Yahyapour et al., 2011; Kahrarian et al., 2011; Daghighi et al., 2013; Darvish-Motevalli, 2016; Hosseini et al., 2016; Ghasemi Charati, 2017; Yahyapour et al., 2020; Ghasemi Charati et al., 2022; Zamani Khormandichali et al., 2024).

**Examined material:** Mazandaran province, Sari, Goharbaran seashore, soil under eucalyptus, 36°42'12"N 53°11'14"E, 5 m a.s.l., 12-May-2020, leg. E. Yahyapour, n= 16.

***Seira dori* Gruia, Poliakov & Broza, 2000**

**General distribution:** Mediterranean (Gruia et al., 2000).

**Distribution in Iran:** Kurdistan (Marivan-Garan) province (Ahmadi et al., 2023).

**Subfamily Willowsiinae Yoshii & Suhardjono, 1989**

**Genus *Drepanosira* Bonet, 1942**

***Drepanosira gisini* Nosek, 1964**

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** South Khorasan (Birjand) province (Rabieh et al., 2023).

***Drepanosira hussi* Neuherz 1976**

**General distribution:** Paleotropical (Baquero et al., 2015).

**Distribution in Iran:** North Khorasan (Darkesh-Jozak forest) province (Mehrafrooz Mayvan et al., 2022).

**Genus *Willowsia* Shoebotham, 1917**

***Willowsia bartkei* Stach, 1965**

**General distribution:** Vietnam (Zhang, 2015).

**Distribution in Iran:** North Khorasan (Darkesh-Jozak forest) province (Mehrafrooz Mayvan et al., 2022).

*Willowsia buski* (Lubbock, 1870)**General distribution:** Cosmopolitan (Bellinger et al., 2024).**Distribution in Iran:** Kogiluyeh and Boyer-Ahmad (Cheram) province (Falahati Hosseinabad et al., 2011).*Willowsia nigromaculata* (Lubbock, 1873)**General distribution:** Holarctic (Babenko et al., 2019).**Distribution in Iran:** Guilan (Rasht) province (Daghighi et al., 2013).**Family Orchesellidae Börner, 1906****Subfamily Heteromurinae Absolon & Kseneman, 1942****Genus *Heteromurtrella* Mari Mutt, 1979***Heteromurtrella* sp.**Distribution in Iran:** North Khorasan (Golil-Sarany rangeland) province (Mehrafrooz Mayvan et al., 2022).**Genus *Heteromurus* Wankel, 1860***Heteromurus gigans* Mari Mutt & Stomp, 1980**General distribution:** Portugal (Arbea, 2021).**Distribution in Iran:** Kurdistan (Marivan-Pirsafa) and Mazandaran (Babol, Sari) provinces (Ghasemi Cherati, 2017; Yahyapour et al., 2020; Ghasemi Charati et al., 2022; Ahmadi et al., 2023).*Heteromurus major* (Moniez, 1889)**General distribution:** Subcosmopolitan (Bellinger et al., 2024).**Distribution in Iran:** Central, East Azarbaijan, Golestan (Gorgan), Guilan (Rasht), Kermanshah (Harsin, Kermanshah, Paveh), Kurdistan (Marivan-Garan, Marivan-Pirsafa), Lorestan, Mazandaran (Babol, Behshahr, Larijan, Neka, Noor, Qaemshahr, Sari, Savadkooh), and Tehran provinces (Cox, 1982; Yahyapour et al., 2011; Daghighi et al., 2013; Qazi & Shayanmehr, 2014; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Balvasi et al., 2015; Hosseini et al., 2016; Mehrafrooz Mayvan et al., 2015; Kahrarian et al., 2014, 2016; Darvish-Motevalli, 2016; Alijani-Ardeshtir et al., 2017; Ghasemi Charati, 2017; Moradi et al., 2018; Bakhshi et al., 2022; Ghasemi Charati et al., 2022; Vahedi Moghadam et al., 2022; Ahmadi et al., 2023; Zamani Khormandichali et al., 2024).**Examined material:** Mazandaran province, Behshahr, Zenget village, Hezarjirib forest, 36°25'15" N, 53°34'16" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n>100; Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n>100; Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 39; Neka, Sapelmiun village, 36°23'11" N, 53°34'18" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 62; Sari, Marzrud village, 36°14'20" N, 53°39'10" E, 46 m a.s.l., 18-May-2020, leg. E. Yahyapour, n>100; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n>100; Sari, Abmal forest, 36°42'52" N, 53°06'00" E, 10 m a.s.l., 24-May-2020, leg. E. Yahyapour, n= 99; Sari, Dashtenaz forest, 36°42'10" N, 53°9'11" E, 11 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 64.*Heteromurus nitidus* (Templeton, 1836) in Templeton & Westwood 1836**General distribution:** Subcosmopolitan (Bellinger et al., 2024).**Distribution in Iran:** Golestan (Kordkouy), Guilan, Kermanshah (Harsin), Kohgiluyeh and Boyer-Ahmad, Mazandaran (Babol, Behshahr, Neka, Qaemshahr, Sari, Savadkooh), and North Khorasan (Darkesh-Jozak forest, Golil-Sarany rangeland, Mayvan village) provinces (Cox, 1982; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Hosseini et al., 2016; Kahrarian, 2015; Kahrarian et al., 2014, 2016; Darvish-Motevalli, 2016; Alijani-Ardeshtir et al., 2017; Ghasemi Cherati,

2017; Bakhshi et al., 2022; Ghasemi Charati et al., 2022; Mehrafrooz Mayvan et al., 2022; Vahedi Moghadam et al., 2022; Zamani Khormandichali et al., 2024).

**Examined material:** Mazandaran province, Behshahr, Zenget village, Hezarjirib forest, 36°25'15" N, 53°34'16" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 4; Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 1; Neka, Sapelmiun village, 36°23'11" N, 53°34'18" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 2; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 9; Sari, Dashtenaz forest, 36°42'10" N, 53°9'11" E, 11 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 1.

*Heteromurus sexoculatus* Brown, 1926

**General distribution:** Asia Minor (Dányi & Traser 2008).

**Distribution in Iran:** Mazandaran and Kermanshah (Char Zabar) provinces (Cox, 1982; Kahrarian et al., 2014; Kahrarian, 2015, 2019).

*Heteromurus variabilis* (Martynova, 1974) in Martynova et al. (1974)

**General distribution:** Tajikistan (Martynova et al., 1974).

**Distribution in Iran:** Kermanshah (Patagh, Sar-e-pole-Zahab), Mazandaran (Behshahr, Sari, Savadkooh) and North Khorasan (Darkesh-Jozak forest) provinces (Kahrarian, 2015; Kahrarian et al., 2015; Ghasemi Charati et al., 2022; Mehrafrooz Mayvan et al., 2022; Vahedi Moghadam et al., 2022; Zamani Khormandichali et al., 2024).

### Subfamily Orchesellinae Börner, 1906

**Genus Orchesella** (Templeton, 1836) in Templeton & Westwood 1836

*Orchesella cincta* (Linnaeus, 1758)

**General distribution:** Subcosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Golestan (Kordkuy) and Mazandaran (Babol, Behshahr, Larijan, Neka, Sari, Savadkooh), North Khorasan (Darkesh-Jozak forest, Golil-Sarany rangeland) provinces (Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Balvasi et al., 2015; Hosseini et al., 2016; Yahyapour et al., 2018, 2020; Ghasemi Charati et al., 2022; Mehrafrooz Mayvan et al., 2022; Vahedi Moghadam et al., 2022).

**Examined material:** Mazandaran province, Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 1; Behshahr, Zenget village, Hezarjirib forest, 36°25'15" N, 53°34'16" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 14; Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 24; Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 14; Neka, Sapelmiun village, 36°23'11" N, 53°34'18" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 1; Sari, Dashtenaz forest, 36°42'10" N, 53°9'11" E, 11 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 1; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 5.

*Orchesella flavescens* (Bourlet, 1839)

**General distribution:** Holarctic (Fjellberg, 2007).

**Distribution in Iran:** North Khorasan (Golil-Sarany rangeland) province (Mehrafrooz Mayvan et al., 2022).

*Orchesella spectabilis* Tullberg, 1871

**General distribution:** Palaearctic (Fjellberg 2007).

**Distribution in Iran:** Mazandaran (Sari) province (Ghasemi Charati, 2017; Ghasemi Charati et al., 2022).

**Family Paronellidae Börner, 1913**  
**Subfamily Cyphoderinae Börner, 1913**  
**Genus *Cyphoderus* Nicolet, 1842**

*Cyphoderus agnotus* Börner, 1906

**General distribution:** Boreal and Neotropical (Lima & Zeppelini, 2021).

**Distribution in Iran:** Golestan (Gorgan, Kordkuy) province (Hosseini et al., 2016).

*Cyphoderus albinus* Nicolet, 1842

**General distribution:** Palaearctic (Fjellberg, 2007).

**Distribution in Iran:** Guilan (Rasht), Isfahan (Zarrinshahr), Kerman (Shahdad), Kermanshah (Parviz Khan, Qasr-e-shirin), Khuzestan, Kohgiluyeh and Boyer-Ahmad, Kurdistan (Marivan-Garan), Mazandaran (Babolsar, Qaemshahr), North Khorasan (Darkesh-Jozak forest), and Tehran provinces (Daghighi, 2012; Qazi & Shayanmehr, 2014; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Darvish-Motevalli, 2016; Kahrarian et al., 2016; Abdolalizadeh, 2018; Kahrarian, 2019; Mehrafrooz Mayvan et al., 2022; Ahmadi et al., 2023).

*Cyphoderus bidenticulatus* Parona, 1888

**General distribution:** Palaearctic (Thibaud 2017).

**Distribution in Iran:** Kermanshah (Harsin, Tamark) and Lorestan provinces (Kahrarian, 2015; Kahrarian et al., 2015; Ghavami, 2016).

**Family Isotomidae Schäffer, 1896**  
**Subfamily Anurophorinae Börner, 1901**  
**Genus *Anurophorus* Nicolet, 1842**

*Anurophorus alpinus* Potapov & Stebaeva, 1990

**General distribution:** Caucasus (Potapov, 2001).

**Distribution in Iran:** Mazandaran (Sari) province (Shayanmehr et al., 2024).

*Anurophorus coiffaiti* Cassagnau & Delamare, 1955

**General distribution:** Armenia, Lebanon, Syria (Potapov, 2001).

**Distribution in Iran:** Golestan (Gorgan), Guilan (Rasht), and Lorestan provinces (Daghighi, 2012; Falahati et al., 2013; Moradi et al., 2018).

*Anurophorus septentrionalis* Palissa, 1966

**General distribution:** Subcosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Mazandaran (Amol, Nowshahr) province (Mohammadi Nodehaki & Shayanmehr, 2020).

*Anurophorus silvaticus* Potapov & Stebaeva, 1990

**General distribution:** Russia and Caucasus (Potapov, 2001).

**Distribution in Iran:** Mazandaran (Behshahr, Qaemshahr, Savadkooh) province (Yoosefi Lafooraki et al., 2019; Vahedi Moghadam et al., 2022).

**Examined material:** Mazandaran province, Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n=3.

**Genus *Bagnallella* Salmon, 1951**

*Bagnallella tenella* (Reuter, 1895)

**General distribution:** Cosmopolitan (Potapov et al., 2021).

**Distribution in Iran:** Guilan (Rasht) province (Daghighi, 2012; Daghighi et al., 2013).

**Genus *Ballistura* Börner, 1906**

*Ballistura schoetti* (Dalla Torre, 1895)

**General distribution:** Cosmopolitan (Potapov, 2001; Fjellberg, 2007).

**Distribution in Iran:** Guilan province (Cox, 1982).

*Ballistura tuberculata* (Stach, 1947)

**General distribution:** Central, Eastern Europe and N America (Potapov, 2001) and Iran

**Distribution in Iran:** Khuzestan province (Ramezani & Mossadegh, 2017).

**Genus *Ephemerotoma* Potapov, Kahrarian, Deharveng & Shayanmehr, 2015**

*Ephemerotoma skarzynskii* Potapov, Kahrarian, Deharveng & Shayanmehr, 2015

**General distribution:** Iran (Potapov et al., 2015).

**Distribution in Iran:** Kermanshah (Cheshmeh Sorkh, Osmanvand) province (Potapov et al., 2015).

**Genus *Folsomia* Willem, 1902**

*Folsomia asiatica* Martynova, 1971

**General distribution:** Tajikistan and Azerbaijan (Potapov, 2001).

**Distribution in Iran:** Golestan (Azadshahr, Kuhmian), Kermanshah (Paveh, Quri Qual'eh), and Lorestan provinces (Amiri & Kahrarian, 2015; Arbea & Kahrarian, 2015a; Ahmadi-Rad & Kahrarian, 2015; Yoosefi Lafooraki et al., 2020b).

*Folsomia binocolata* (Wahlgren, 1899)

**General distribution:** Palaearctic (Fjellberg, 2007).

**Distribution in Iran:** Kermanshah province (Ghahramaninezhad et al., 2013).

*Folsomia brevifurca* (Bagnall, 1949)

**General distribution:** Palaearctic (Dányi and Traser, 2008).

**Distribution in Iran:** Mazandaran province (Cox, 1982).

*Folsomia candida* (Willem, 1902)

**General distribution:** Cosmopolitan (Potapov, 2001; Fjellberg, 2007).

**Distribution in Iran:** Central, East Azarbaijan, Golestan (Aliabad, Azadshahr, Kuhmian), Guilan (Fuman, Rezwanshahr), Kurdistan (Marivan-Pirsafa), and Mazandaran (Abbas-Abad, Sari) provinces (Cox, 1982; Yahyapour, 2012; Yoosefi Lafooraki et al., 2020b; Ahmadi et al., 2023).

*Folsomia diplophthalma* (Axelson, 1902)

**General distribution:** Palaearctic (Babenko et al., 2019).

**Distribution in Iran:** Kerman (Mahan) province (Abdolalizadeh, 2018; Abdolalizadeh et al., 2018).

*Folsomia fimetaria* (Linnaeus, 1758)

**General distribution:** Holarctic (Babenko et al., 2019).

**Distribution in Iran:** Central, Guilan, and Mazandaran provinces (Cox, 1982).

*Folsomia inoculata* Stach, 1947

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** Mazandaran (Sari) province (Yahyapour et al., 2020).

*Folsomia ksenemani* Stach, 1947

**General distribution:** Europe, West Caucasus, Lebanon (Potapov, 2001).

**Distribution in Iran:** Mazandaran (Noor, Savadkooh) and Tehran provinces (Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Qazi and Shayanmehr, 2016).

*Folsomia manolachei* Bagnall, 1939

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** Golestan (Gorgan), Kermanshah, Kurdistan (Marivan-Pirsafa), and Mazandaran (Behshahr) provinces (Arbea & Kahrarian, 2015b; Yoosefi Lafooraki et al., 2020b; Vahedi Moghadam et al., 2022; Ahmadi et al., 2023).

**Examined material:** Mazandaran province, Behshahr, Zenget village, Hezarjirib forest, 36°25'15" N, 53°34'16" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 5; Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 7; Neka, Sapelmiun village, 36°23'11" N, 53°34'18" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 20.

*Folsomia penicula* Bagnall, 1939

**General distribution:** Holarctic (Bellinger et al., 2024).

**Distribution in Iran:** Central, Guilan (Astaneh-ashrafiyeh, Fuman, Rasht, Rezwanshahr, Rudbar, Shaft), Kermanshah, Kurdistan (Marivan-Pirsafa), Mazandaran (Abbas-Abad, Amol, Behshahr, Chalus, Galugah, Neka, Noor, Nowshahr, Qaemshahr, Sari, Savadkooh, Tonekabon), East Azarbaijan, Golestan (Aliabad, Bandar Gaz), North Khorasan (Darkesh-Jozak forest), and Tehran provinces (Cox, 1982; Falahati et al., 2011; Daghighi, 2012; Falahati et al., 2013; Daghighi et al., 2013; Qazi & Shayanmehr, 2014; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Arbea & Kahrarian, 2015; Yoosefi Lafooraki et al., 2020b; Bakhshi et al., 2022; Ghasemi Charati et al., 2022; Mehrafrooz Mayvan et al., 2022; Ahmadi et al., 2023; Zamani Khormandichali et al., 2024).

**Examined material:** Mazandaran Province, Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 7; Neka, Sapelmiun village, 36°23'11" N, 53°34'18" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 9; Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 1; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 21.

*Folsomia quadrioculata* (Tullberg, 1871)

**General distribution:** Subcosmopolitan (Checklist of the Collembola of the World 2024).

**Distribution in Iran:** Central, East Azarbaijan, Golestan (Bandar Gaz, Nowkandeh), Guilan (Shaft), Ilam (Choqa-Sabz Forest Park), Kermanshah (Char zabar, Shikh salaeh village, Ghap Gholi village, Sia Khor village), Kurdistan (Marivan-Pirsafa), Lorestan, Mazandaran (Abbas-Abad, Nowshahr), North Khorasan (Darkesh-Jozak forest, Golil-Sarany rangeland), and West Azarbaijan provinces (Cox, 1982; Amiri & Kahrarian, 2015; Arbea & Kahrarian, 2015a; Moradi et al., 2018; Shayanmehr et al., 2019; Yoosefi Lafooraki et al., 2020b; Mehrafrooz Mayvan et al., 2022; Ahmadi et al., 2023).

*Folsomia similis* Bagnall, 1939

**General distribution:** Holarctic (Babenko et al., 2019)

**Distribution in Iran:** Guilan (Rasht), Kermanshah (Paveh), and Tehran provinces (Moravvej et al., 2007; Daghighi, 2012; Amiri & Kahrarian, 2015; Arbea & Kahrarian, 2015a).

*Folsomia trisetata* Jordana & Ardanaz, 1981

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** Mazandaran (Neka) province (Bakhshi et al., 2022).

**Genus *Folsomides* Stach, 1922**

***Folsomides angularis*** (Axelson, 1905)

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** Guilan (Rasht, Rudbar, Shaft) and Kurdistan (Marivan-Pirsafa) provinces (Daghighi et al., 2013; Yoosefi Lafooraki et al., 2020b; Ahmadi et al., 2023).

***Folsomides halshinicus*** Arbea & Kahrarian, 2015

**General distribution:** Iran (Arbea & Kahrarian, 2015a).

**Distribution in Iran:** Kermanshah (Char zabar, Hassan Abad village, Islam Abad-e Gharb, Shabankareh village, Sia Khor village) and Kurdistan (Marivan-Pirsafa) provinces (Kahrarian, 2015; Ahmadi et al., 2023).

***Folsomides marchicus*** (Frenzel, 1941)

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** Ilam (Choqa-Sabz Forest Park, Dalab Forest Park, Gachan Mountains), Kermanshah (Eslamabade-Gharb), Kurdistan (Marivan-Garan, Marivan-Pirsafa), and Lorestan provinces (Kahrarian et al., 2012; Amiri & Kahrarian, 2015; Arbea & Kahrarian, 2015a; Kahrarian, 2015, 2019; Moradi et al., 2018; Shayanmehr et al., 2019; Ahmadi et al., 2023).

***Folsomides parvulus*** (Starch, 1922)

**General distribution:** Cosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Central, East Azarbaijan, Golestan (Gonbade Qabus, Kordkuy, Minudasht), Guilan (Rasht), Kerman (Mahan), Kermanshah (Chahar Zabar-e-Oliya, Harsin), Kurdistan (Marivan-Garan), Lorestan, Mazandaran (Babol, Babolsar, Neka, Qaemshah, Sari, Savadkooh), North Khorasan (Golil-Sarany rangeland), Semnan (Mahdishahr), Tehran, and West Azarbaijan provinces (Cox, 1982; Yahyapour, 2012; Kahrarian et al., 2012; Daghighi, 2012; Daghighi et al., 2013; Qazi & Shayanmehr, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Amiri & Kahrarian, 2015; Kahrarian, 2015; Hosseini et al., 2016; Alijani-Ardeshtir et al., 2017; Arbea & Kahrarian, 2017; Abdolalizadeh, 2018; Moradi et al., 2018; Yoosefi Lafooraki et al., 2020b; Bakhshi et al., 2022; Mehrafrooz Mayvan et al., 2022; Ahmadi et al., 2023; Zamani Khormandichali et al., 2024).

**Examined material:** Mazandaran province, Behshahr, Zenget village, Hezarjirib forest, 36°25'15" N, 53°34'16" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 1.

***Folsomides subvinosus*** Arbea & Kahrarian, 2015

**General distribution:** Iran (Arbea and Kahrarian, 2015a,b).

**Distribution in Iran:** Kermanshah (Char zabar, Islam Abad-e Gharb, Shabankareh village, Sia Khor village) province (Arbea & Kahrarian, 2015; Kahrarian, 2015).

**Genus *Gnathofolsomia* Deharveng & Christian, 1984**

***Gnathofolsomia* sp.**

**Distribution in Iran:** Kermanshah province (Shayanmehr et al., 2019, 2020).

**Genus *Hemisotoma* Bagnall, 1949**

***Hemisotoma orientalis*** (Stach, 1947)

**General distribution:** Ukraine, Kazakhstan, S Siberia, SE France (Potapov, 2001).

**Distribution in Iran:** Central, East Azarbaijan, Guilan, Kermanshah (Chahar Zebar-e-Oliya, Patagh), Mazandaran, and West Azarbaijan provinces (Cox, 1982; Amiri & Kahrarian, 2015; Arbea & Kahrarian, 2015a; Kahrarian, 2015).

***Hemisotoma pontica*** Stach, 1947

**General distribution:** Palaearctic (Bellinger et al., 2024)

**Distribution in Iran:** Central, East Azarbaijan, Golestan (Aqqala, Bandar Torkmen), Guilan, Ilam (Arghavan Forest Park), Kermanshah, Kurdistan (Marivan-Garan, Marivan-Pirsafa), Lorestan, Mazandaran (Amol, Babol, Mahmood-abad, Qaemshahr, Sari, Simorq), North Khorasan (Darkesh-Jozak forest), Tehran, and West Azarbaijan provinces (Cox, 1982; Moravvej et al., 2007; Kahrarian et al., 2012; Yahyapour, 2012; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Darvish-Motevalli, 2016; Alijani-Ardeshtir et al., 2017; Moradi et al., 2018; Shayanmehr et al., 2019; Yoosefi Lafooraki et al., 2020b; Mehrafrooz Mayvan et al., 2022; Ahmadi et al., 2023).

***Hemisotoma quadrioculata*** (Martynova, 1968): Martynova, 1968a

**General distribution:** Middle Asia, Tajikistan (Potapov, 2001).

**Distribution in Iran:** Ilam (Arghavan Forest Park) and Kurdistan (Marivan-Pirsafa) provinces (Shayanmehr et al., 2017; Ahmadi et al., 2023).

***Hemisotoma thermophila*** (Axelson, 1900)

**General distribution:** Cosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Central, East Azarbaijan, Golestan (Aqqala, Azadshahr, Galikesh, Gorgan, Kordkuy, Maraveh Tappeh, Minudasht), Guilan (Rasht, Masal), Kermanshah (Derkah village, Shikh salaeh village, Rijab), Mazandaran (Amol, Larijan, Mahmood-abad, Noor), and West Azarbaijan provinces (Cox, 1982; Daghighi, 2012; Daghighi et al., 2013; Yoosefi Lafooraki (2014), Yoosefi Lafooraki & Shayanmehr, 2014a; Balvasi et al., 2015; Arbea & Kahrarian, 2015a; Yoosefi Lafooraki et al., 2020b).

**Examined material:** Mazandaran province, Sari, Goharbaran seashore, soil under eucalyptus, 36°42'12" N 53°11'14" E, 5 m a.s.l., 12-May-2020, leg. E. Yahyapour, n= 1.

**Genus *Isotomiella*** Bagnall, 1939

***Isotomiella gracilimucronata*** Rusek, 1981

**General distribution:** Iraq (Potapov, 2001).

**Distribution in Iran:** North Khorasan (Golil-Sarany rangeland) province (Mehrafrooz Mayvan et al., 2022).

***Isotomiella minor*** (Schäffer, 1896)

**General distribution:** Cosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** East Azarbaijan, Golestan (Ali-abad, Bandar Gaz, Gorgan, Kordkuy, Ramian), Guilan (Fuman, Lahijan, Rasht, Rudsar, Talesh), Kerman (Koohpayeh village), Kermanshah (Shabankareh village, Patagh), Kurdistan (Marivan-Garan), Lorestan, Mazandaran (Abbas-Abad, Amol, Behshahr, Qaemshahr, Neka, Noor, Nowshahr, Sari, Savadkooh), North Khorasan (Golil-Sarany rangeland), and Tehran provinces (Cox, 1982; Moravvej et al., 2007; Yahyapour, 2012; Daghighi, 2012; Ghahramaninezhad et al., 2013; Daghighi et al., 2013; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Amiri & Kahrarian, 2015; Arbea & Kahrarian 2015a; Kahrarian, 2015, 2019; Hosseini et al., 2016; Ghasemi Charati, 2017; Abdolalizadeh, 2018; Moradi et al., 2018; Yoosefi Lafooraki et al., 2020b; Bakhshi et al., 2022; Ghasemi Charati et al., 2022; Mehrafrooz Mayvan et al., 2022; Ahmadi et al., 2023; Zamani Khormandichali et al., 2024).

**Examined material:** Mazandaran Province, Sari, Dashtenaz forest, 36°42'10" N, 53°9'11" E, 11 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 1.

**Genus *Proisotoma* Börner, 1903**

***Proisotoma minima* Absolon, 1901**

**General distribution:** Holarctic with some records from the southern hemisphere (Babenko et al. 2019).

**Distribution in Iran:** Golestan (Golestan National Park) and Mazandaran (Sari) provinces (Yoosefi Lafooraki & Shayanmehr, 2014; Yahyapour et al., 2020).

***Proisotoma minuta* (Tullberg, 1871)**

**General distribution:** Cosmopolitan (Babenko et al., 2019).

**Distribution in Iran:** Central, East Azarbaijan, Golestan (Kordkuy, Palangpa forest), Guilan (Parehsar, Rasht, Rezwanshahr, Rudbar), Isfahan, Mazandaran (Chalus, Larijan, Mahmoodabad), and West Azarbaijan provinces (Cox, 1982; Nematollahi et al., 2009; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Balvasi et al., 2015; Hosseini et al., 2016; Yoosefi Lafooraki et al., 2020b).

***Proisotoma subminuta* Denis, 1931**

**General distribution:** Cosmopolitan (Fjellberg, 2007).

**Distribution in Iran:** Golestan, Guilan (Rasht), Mazandaran (Amol), and Tehran provinces (Moravvej et al., 2007; Daghighi, 2012; Daghighi et al., 2013; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Yoosefi Lafooraki et al., 2020b).

**Genus *Subisotoma* (Stach, 1947)**

***Subisotoma variabilis* (Gisin, 1949)**

**General distribution:** Switzerland, Austria, Poland, Slovakia, France, Iberian Peninsula, Siberia, Kazakhstan, and Central Asia (Potapov, 2001).

**Distribution in Iran:** Kermanshah (Patagh, Sar-e-pole-Zahab) province (Kahrarian, 2015; Kahrarian et al., 2015).

**Subfamily Isotominae Schäffer, 1896**

**Genus *Agrenia* Börner, 1906**

***Agrenia* sp.**

**Distribution in Iran:** North Khorasan (Zoeram village) province (Mehrafrooz Mayvan et al., 2022).

**Genus *Desoria* Agassiz & Nicolet, 1841 in Desor (1841)**

***Desoria neglecta* (Schäffer, 1900)**

**General distribution:** Holarctic (Fjellberg, 2007).

**Distribution in Iran:** Kurdistan (Marivan-Pirsafa), Lorestan, and Mazandaran (Savadkooh) provinces (Moradi et al., 2018; Ahmadi et al., 2023; Zamani Khormandichali et al., 2024).

***Desoria kaszabi* Dunger, 1982\***

**General distribution:** Palaearctic (Bellinger et al., 2024).

**Distribution in Iran:** This species is recorded for the first time from Iran.

**Examined material:** Mazandaran province, Sari, Marzrud village, 36°14'20" N, 53°39'10" E, 46 m a.s.l., 18-May-2020, leg. E. Yahyapour, n=47; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l.,

20-May-2020, leg. E. Yahyapour, n= 4; Sari, Dashtenaz forest, 36°42'10" N, 53°9'11" E, 11 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 2.

***Desoria olivacea*** (Tullberg, 1871)

**General distribution:** Holarctic (Fjellberg, 2007).

**Distribution in Iran:** East Azarbaijan province (Cox, 1982).

***Desoria tigrina*** Nicolet, 1842

**General distribution:** Palaearctic (Fjellberg, 2007).

**Distribution in Iran:** Kermanshah (Derkah) and Lorestan provinces (Kahrarian et al., 2012; Kahrarian, 2015; Kahrarian & Arbea, 2015; Arbea & Kahrarian, 2015a; Kahrarian, 2019; Moradi et al., 2018).

***Desoria trispinata*** (MacGillivray, 1896)

**General distribution:** Cosmopolitan (Babenko et al., 2019).

**Distribution in Iran:** North Khorasan (Darkesh-Jozak forest) province (Mehrafrouz Mayvan et al., 2022).

***Desoria zlotini*** Martynova, 1968: Martynova, 1968b

**General distribution:** Kirghisia, Tajikistan, Mongolia (Potapov, 2001).

**Distribution in Iran:** Kermanshah (Osmanevad, Sragool village) province (Amiri & Kahrarian, 2015; Kahrarian et al., 2015).

**Genus *Isotoma*** Bourlet, 1839

***Isotoma anglicana*** Lubbock, 1862

**General distribution:** Subcosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Ilam (Arghavan Forest Park, Dalab Forest Park, Gachan Mountain) province (Shaynmehr et al., 2017).

Additional localities: Mazandaran province, Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 2; Sari, Dashtenaz forest, 36°42'10" N, 53°9'11" E, 11 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 2.

***Isotoma iranica*** Arbea & Kahrarian, 2015

**General distribution:** Iran (Arbea and Kahrarian, 2015b).

**Distribution in Iran:** Golestan (Galikesh, Minudasht, Ramian), Kermanshah (Paveh, Shahu city), Lorestan, and Mazandaran (Savadkooh) provinces (Arbea & Kahrarian, 2015a; Moradi et al., 2018; Yoosefi Lafooraki et al., 2020b; Zamani Khormandichali et al., 2024).

Additional localities: Mazandaran province, Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 1.

***Isotoma cf. riparia*** (Nicolet, 1842)

**General distribution:** Palaearctic (Fjellberg, 2007).

**Distribution in Iran:** Mazandaran (Babol) province (Alijani-Ardeshtir et al., 2017).

***Isotoma viridis*** Bourlet, 1839

**General distribution:** Holarctic (Babenko et al., 2019).

**Distribution in Iran:** Central, East Azarbaijan, Golestan (Golestan National Park), Kermanshah (Paveh, Shabankareh), Kurdistan (Marivan-Garan, Marivan-Pirsafa), Mazandaran (Babol, Sari), North Khorasan (Darkesh-Jozak forest), and West Azarbaijan provinces (Cox, 1982; Yahyapour, 2012; Amiri & Kahrarian, 2015; Kahrarian, 2015, 2019; Alijani-Ardeshtir et al., 2017; Mehrafrouz Mayvan et al., 2022; Ahmadi et al., 2023).

**Genus *Isotomodella* Martynova, 1968*****Isotomodella alticola*** (Bagnall, 1949)**General distribution:** Palaearctic (Fjellberg, 2007).**Distribution in Iran:** Lorestan (Khoram-Abad, Veysian, Delbar Sadat village) province (Shayanmehr et al., 2018).**Genus *Isotomodes* Axelson, 1907*****Isotomodes golestanii*** Yoosefi Lafooraki, Hajizadeh, Shayanmehr & Hosseini, 2020**General distribution:** Iran (Yoosefi Lafooraki et al., 2020b).**Distribution in Iran:** Golestan (Gonbade Qabus) province (Yoosefi Lafooraki et al., 2020b).***Isotomodes korkorensis*** Arbea & Kahrarian, 2015**General distribution:** Iran (Arbea & Kahrarian, 2015b).**Distribution in Iran:** Kermanshah (Char zabar, Kor Kor Mountain) province (Arbea & Kahrarian, 2015a).**Genus *Isotomurus* Börner, 1903*****Isotomurus afghanicus*** Yosii, 1966**General distribution:** Iran and Afghanistan (Yoosefi Lafooraki et al., 2023).**Distribution in Iran:** Mazandaran (Amol, Mahmood-Abad, Neka, Noor, Nowshahr, Sari) province (Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Bakhshi et al., 2022; Yoosefi Lafooraki et al., 2023).**Additional localities:** Mazandaran province, Sari, Marzrud village, 36°14'20" N, 53°39'10" E, 46 m a.s.l., 18-May-2020, leg. E. Yahyapour, n= 76; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 14; Sari, Abmal forest, 36°42'52" N, 53°06'00" E, 10 m a.s.l., 24-May-2020, leg. E. Yahyapour, n= 13; Sari, Dashtenaz forest, 36°42'10" N, 53°09'11" E, 11 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 6.***Isotomurus antennalis*** Bagnall, 1940**General distribution:** Palaearctic (Potapov, 2001).**Distribution in Iran:** Mazandaran (Babol) province (Alijani-Ardeshtir et al., 2017).***Isotomurus cf. balteatus*** (Reuter, 1876)**General distribution:** Unclear (revision needed) (Fjellberg, 2007).**Distribution in Iran:** Golestan (Kordkuy) province (Hosseini et al., 2016).***Isotomurus cassagnau*** Deharveg & Lek, 1993**General distribution:** France (Thibaud, 2017).**Distribution in Iran:** Mazandaran (Sari) province (Zamani, 2016).***Isotomurus fucicola*** (Schött, 1893)**General distribution:** Subcosmopolitan (Bellinger et al., 2024).**Distribution in Iran:** Mazandaran (Qaemshahr) province (Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014).**Additional localities:** Mazandaran province, Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 3; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 1.

*Isotomurus graminis* Fjellberg, 2007**General distribution:** Palaearctic (Bellinger et al., 2024).**Distribution in Iran:** North Khorasan (Golil-Sarany rangeland) province (Mehrafrooz Mayvan & Greenslade, 2020; Mehrafrooz Mayvan et al., 2022).*Isotomurus hyrcanicus* Yoosefi Lafooraki & Shayanmehr, 2023 in Yoosefi Lafooraki et al. (2023)**General distribution:** Iran (Yoosefi Lafooraki et al., 2023).**Distribution in Iran:** Golestan (Bandar Gaz, Bandar Torkaman) and Mazandaran (Sari, Savadkooh) provinces (Yoosefi Lafooraki et al., 2023; Zamani Khormandichali et al., 2024).*Isotomurus katule* Yoosefi Lafooraki & Shayanmehr, 2023 in Yoosefi Lafooraki et al. (2023)**General distribution:** Iran (Yoosefi Lafooraki et al., 2023).**Distribution in Iran:** Golestan (Aliabad-e katul) and Mazandaran (Savadkooh) provinces (Yoosefi Lafooraki et al., 2023; Zamani Khormandichali et al., 2024).*Isotomurus maculatus* (Schäffer, 1896)**General distribution:** unclear (revision needed) (Fjellberg, 2007).**Distribution in Iran:** Golestan (Gorgan) and Mazandaran (Sari) provinces (Falahati et al., 2011, 2013; Yoosefi Lafooraki et al., 2023).*Isotomurus matanicus* Shayanmehr, Yoosefi & Zamani, 2024 in Zamani Khormandichali et al. 2024**General distribution:** Iran (Zamani Khormandichali et al., 2024).**Distribution in Iran:** Mazandaran (Savadkooh) province (Zamani Khormandichali et al., 2024).*Isotomurus palustris* (Müller, 1776)**General distribution:** Cosmopolitan (Bellinger et al., 2024).**Distribution in Iran:** Central, Golestan (Kordkuy, Palangpa forest, Imam Reza forest), Guilan, Khuzestan, and Mazandaran (Sari) provinces (Cox, 1982; Hosseini et al., 2016; Zamani, 2016; Ramezani & Mossadegh, 2017).

Additional localities: Mazandaran province, Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 2; Sari, Marzrud village, 36°14'20" N, 53°39'10" E, 46 m a.s.l., 18-May-2020, leg. E. Yahyapour, n= 2; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 1.

*Isotomurus potapovi* Yoosefi Lafooraki & Shayanmehr, 2023 in Yoosefi Lafooraki et al. (2023)**General distribution:** Iran (Yoosefi Lafooraki et al., 2023).**Distribution in Iran:** Golestan (Bandar Gaz, Kordkuy), Guilan (Amlash, Falakdeh, Ostad-Kalayeh, Rasht, Roodbar), and Mazandaran (Abbas-Abad, Savadkooh, Tonekabon) provinces (Yoosefi Lafooraki et al., 2023; Zamani Khormandichali et al., 2024).*Isotomurus punctiferus* Yosii, 1963**General distribution:** Palaearctic (Bellinger et al., 2024).**Distribution in Iran:** Golestan (Gorgan) and Guilan (Rasht) provinces (Falahati et al., 2013; Daghighi et al., 2013; Yoosefi Lafooraki et al., 2023).

Additional localities: Mazandaran province, Sari, rice field, 36°42'10" N, 53°10'12" E, 11 m a.s.l., 12-May-2020, leg. E. Yahyapour, n= 30.

*Isotomurus stuxbergi* (Tullberg, 1876)**General distribution:** Holoarctic (Babenko et al., 2019).**Distribution in Iran:** Mazandaran (Qaemshahr) province (Darvish-Motevalli, 2016).

**Genus *Pachyotoma* Bagnall, 1949*****Pachyotoma* sp.**

**Distribution in Iran:** Kermanshah province (Amiri & Kahrarian, 2015).

**Genus *Parisotoma* Bagnall, 1940*****Parisotoma notabilis* (Schäffer, 1896)**

**General distribution:** Cosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Central, East Azarbaijan, Golestan (Aliabd, Bandar Gaz, Bandar Torkmen, Golestan National Park, Kalaleh, Kordkuy, Maraveh Tappe, Palangpa forest), Guilan (Astaneh-ashrafiyeh, Fuman, Langrud, Rasht, Rezwanshahr, Rudbar, Rudsar, Shaft, Somesara, Talesh), Kerman (Mahan, Shahdad), Kermanshah (Patagh, Shabankareh villag), Kermanshah, Khuzestan, Kurdistan (Marivan-Garan, Marivan-Pirsafa), Lorestan, Mazandaran (Behshahr, Chalus, Galugah, Qaemshahr, Kodirsar, Mahmood-Abad, Neka, Noor, Nowshahr, Royan, Savadkooh), North Khorasan (Darkesh-Jozak forest, Gelyan village, Golil-Sarany rangeland), Tehran, West Azarbaijan, and Zanzan provinces (Cox, 1982; Moravvej et al., 2007; Kahrarian et al., 2012; Daghighi et al., 2013; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Amiri & Kahrarian, 2015; Arbea & Kahrarian, 2015a; Kahrarian, 2015, 2019; Hosseini et al., 2016; Ramezani & Mossadegh, 2017; Abdolalizadeh, 2018; Moradi et al., 2018; Yoosefi Lafooraki et al., 2020; Bakhshi et al., 2022; Mehrafrooz Mayvan et al., 2022; Ahmadi et al., 2023; Zamani Khormandichali et al., 2024).

Additional localities: Mazandaran province, Behshahr, Zenget village, Hezarjirib forest, 36°25'15" N, 53°34'16" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 3; Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 46; Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 16; Neka, Sapelmiun village, 36°23'11" N, 53°34'18" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 15; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 56; Sari, Abmal forest, 36°42'52" N, 53°06'00" E, 10 m a.s.l., 24-May-2020, leg. E. Yahyapour, n= 3; Sari, Dashtenaz forest, 36°42'10" N, 53°9'11" E, 11 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 13.

**Genus *Pseudisotoma* Handschin, 1924*****Pseudisotoma sensibilis* (Tullberg, 1876)**

**General distribution:** Palaearctic (Fjellberg, 2007).

**Distribution in Iran:** Kurdistan (Marivan-Garan, Marivan-Pirsafa) and Mazandaran (Noor, Royan) provinces (Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Ahmadi et al., 2023).

**Genus *Vertagopus* Börner, 1906*****Vertagopus persicus* Potapov, Yoosefi & Shayanmehr, 2020 in Yoosefi Lafooraki et al. 2020a**

**General distribution:** West Asia (Bellinger et al., 2024).

**Distribution in Iran:** Golestan (Nokandeh), Guilan (Shaft), and Mazandaran (Abbas-Abad) provinces (Yoosefi Lafooraki et al., 2020a).

Additional localities: Mazandaran Province, Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 24; Neka, Sapelmiun village, 36°23'11" N, 53°34'18" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 2; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 10.


**Family Oncopoduridae Carl & Lebedinsky, 1905****Genus *Oncopodura* Carl & Lebedinsky, 1905***Oncopodura ambigua* Christiansen, 1957**General distribution:** Lebanon (Christiansen, 1957).**Distribution in Iran:** Guilan and Mazandaran provinces (Cox, 1982).*Oncopodura hamata* Carl & Lebedinsky, 1905**General distribution:** Europe (Arbea, 2021).**Distribution in Iran:** Guilan province (Daghighi, 2012).**Family Tomoceridae Schäffer, 1896****Genus *Pogonognathellus* Paclt, 1944***Pogonognathellus flavescens* (Tullberg, 1871) (Figure 2)**General distribution:** Palaearctic and Nearctic (Babenko et al., 2019).**Distribution in Iran:** Mazandaran (Babol, Neka, Sari, Tonekabon) province (Yahyapour et al., 2018).**Genus *Tomocerina* Yosii, 1955***Tomocerina minuta* (Tullberg, 1876) (Figure 3B)**General distribution:** Subcosmopolitan (Bellinger et al., 2024).**Distribution in Iran:** Mazandaran (Neka, Sari) province (Yahyapour et al., 2021; Bakhshi et al., 2022).**Examined material:** Mazandaran province, Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 8; Behshahr, Zenget village, Hezarjirib forest, 36°25'15" N, 53°34'16" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 1; Neka, Lemon village, Hezarjirib forest, 36°27'18" N, 53°34'04" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 10; Neka, Sapelmiun village, 36°23'11" N, 53°34'18" E, 1057 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 8; Sari, Dashtenaz forest, 36°42'10" N, 53°9'11" E, 11 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 60; Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 17.**Genus *Tomocerus* Nicolet, 1842***Tomocerus asiaticus* Martynova 1969\***General distribution:** West and Central Asia (Bellinger et al., 2024).**Distribution in Iran:** The species is reported for the first time from Iran.

Additional localities: Mazandaran province, Sari, Zarrin-Abad forest, 36°32'44" N, 53°07'53" E, 113 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 1; Sari, Dashtenaz forest, 36°42'10" N, 53°9'11" E, 11 m a.s.l., 20-May-2020, leg. E. Yahyapour, n= 1.

*Tomocerus minor* (Lubbock, 1862) (Figure 4A)**General distribution:** Subcosmopolitan (Bellinger et al., 2024).**Distribution in Iran:** Central, East Azarbaijan, Golestan, Guilan, Kerman, Mazandaran, and Zanjan provinces (Cox, 1982; Hosseini et al., 2016; Abdolalizadeh, 2018).**Examined material:** Mazandaran province, Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 4.



**FIGURE 2.** Inner side of dens with a broad pointed scale near base base in *Pogonognathellus flavescens* (Yahyapour et al., 2018).




**FIGURE 3.** A) Mucro with two dorsal lamella, mid-section of mucro with more than three teeth in *Tomocerus vulgaris*, B) Mid-section of mucro at most with three teeth in *Tomocerina minuta* (Magnification 40X).

***Tomocerus vulgaris*** (Tullberg, 1871) (Figure 3B, 4B)

**General distribution:** Subcosmopolitan (Bellinger et al., 2024).

**Distribution in Iran:** Central, East Azarbaijan, Golestan, Guilan, Kermanshah, Mazandaran (Behshahr, Sari, Tonekabon), North Khorasan (Darkesh-Jozak forest), and Zanzan provinces (Cox, 1982; Yoosefi Lafooraki, 2014; Yoosefi Lafooraki & Shayanmehr, 2014; Balvasi et al., 2015; Kahrarian, 2015; Kahrarian et al., 2014, 2015; Hosseini et al., 2016; Alijani-Ardeshtir et al., 2017; Yahyapour et al., 2020; Ghasemi Charati et al., 2022; Mehrafrooz Mayvan et al., 2022; Vahedi Moghadam et al., 2022).

**Examined material:** Mazandaran Province, Behshahr, Rudbar village, Hezarjirib forest, 36°27'18" N, 53°34'47" E, 46 m a.s.l., 10-May-2020, leg. E. Yahyapour, n= 1; Sari, Abmal forest, 36°42'52" N, 53°06'00" E, 10 m a.s.l., 24-May-2020, leg. E. Yahyapour, n= 1.



**FIGURE 4.** A) Dorsal spines of dens 3-forked in *Tomocerus minor*, B) Simple spines on dorsal side of dens in *Tomocerus vulgaris* (Magnification 100X).

**Key to species of Tomoceridae family in Iran (according to Fjellberg, 2007):**

- 1- Inner side of dens with a broad pointed scale near base (Fig. 3). Basolateral field of labium with more than five setae. Mucro with one dorsal lamella ..... *ogonognathellus flavescens* (Tullberg, 1871)
- Inner side of dens without broad scale. Basolateral field of labium with 5 setae. Mucro with two dorsal lamella (Fig. 4), the outer one with small teeth ..... 2
- 2- All tibiotarsi with several differentiated macrochaetae on inner side. Mid-section of mucro with more than 3 teeth (Fig. 4), outer basal tooth with a small secondary tooth .... *Tomocerus* Nicolet 3
- Tibiotarsi 1-2 without, tib. 3 with one inner macrochaetae. Mid-section of mucro at most with 3 teeth (Fig. 5), outer basal tooth simple ..... *Tomocerina minuta* (Tullberg, 1876)
- 3- Dorsal spines of dens 3-forked (Fig. 6) ..... *Tomocerus minor*
- Dorsal spines of dens simple (Fig. 7) ..... 4
- 4- Empodium without teeth ..... *Tomocerus asiaticus*
- Empodium with one inner tooth ..... *Tomocerus vulgaris*

**DISCUSSION**

Our comprehensive study documents 139 species from 41 genera across six families of Entomobryomorpha in Iran, including three newly reported species. The family-level diversity shows Isotomidae as the most speciose (69 species, 21 genera), followed by Entomobryidae (51 species, 12 genera), Orchesellidae (9 species, 3 genera), Tomoceridae (5 species, 3 genera), Paronellidae (3 species, 1 genus), and Oncopoduridae (2 species, 1 genus). A particularly noteworthy finding emerges when comparing our results with the previous checklist by Shayanmehr et al. (2020). In just five years, we've witnessed remarkable growth in documented Entomobryomorpha diversity: from 34 to 105 species (309% increase) in the Isotomidae family, from 34 total species reported in 2020 to 139 species currently known, 2.9-fold increase in recorded species richness, and 15 new genera added to Iran's fauna. This dramatic expansion can be attributed to intensive sampling in previously unexplored regions and re-examination of historical collections.

The rapid discovery rate suggests Iran likely represents a biodiversity hotspot for Collembola and many species remain to be discovered. Therefore we recommend targeted sampling in understudied ecoregions and development of regional identification keys. This study establishes a new baseline for understanding Collembola diversity in Western Asia and highlights the importance of sustained taxonomic research on microarthropods.

### Acknowledgments

This work has been supported by the Vice Chancellor for Research in the Sari University of Agricultural Sciences and Natural Resources (SANRU) for financing the research project with contract 32 number 01-1398-01. Also, the authors would like to thank and the anonymous reviewers for their valuable suggestions and corrections that improved this paper.

### LITERATURE CITED

Abdolalizadeh, F. 2018. A survey on Collembola fauna in Kerman County. MsC thesis, Shahid Bahonar University of Kerman, 90p.

Abdolalizadeh, F., Khayrandish, M., Shayanmehr, M., Bernard C.E., 2019. Introduction of springtails (Collembola) from Kerman city and its suburbs with some new records for Iran fauna. *Entomology and phytopathology journal* 87(1), 63–75. (in Persian with English abstract) <https://doi.org/10.22092/jaep.2019.123392.1247>

Absolon, K., 1901. Zwei neue Collembolenformen aus den Höhlen des mährischen Karstes. *Zoologischer Anzeiger* 24(634), 32–33.

Absolon, K., Kseneman, M., 1942. Troglopedetini. Vergleichende Studie über eine altertümliche höhlenbewohnende Kollembolengruppe aus den dinarischen Karstgebieten. *Studien aus dem Gebiete der allgemeinen Karstforschung, der wissenschaftlichen Höhlenkunde, der Eiszeitforschung und den Nachbargebieten.*, Biologische Serie (Nr. 16), Brünn 5–57.

Ahmadi, S., Ghobari, H., Shayanmehr, M., Mohammadi-Samani, K., Kapruś, I., 2023. Addition to Iranian Springtails fauna and a checklist of the Collembola from Kurdistan province. *Journal of Insect Biodiversity and Systematics* 9(1), 01–16. <https://doi.org/10.52547/jibs.9.1.1>

Ahmadi-Rad, A., Kahrarian, M. 2015. The first study of Isotomidae fauna (Collembola: apterygota) in Lorestan Province, Western Iran, with new records for Iranian fauna. *Journal of Biodiversity and Environmental Sciences* 7(1), 128–133.

Alijani-Ardeshtir, M., Shayanmehr, M., Amiri-Besheli, B. 2017. Faunistic survey of Collembola in Babol regions, Mazandaran province: Introduction new species for Iranian fauna. *Plant Protection (Scientific Journal of Agriculture)* 40(2), 45–61 [In Persian with English summary].

Amiri, H., Kahrarian, M. 2015. The new record of Isotomidae (Collembola: Apterygota) for Iranian Fauna: an update to the species Listin Kermanshah Province (Western Iran). *Journal of Biodiversity and Environmental Sciences (JBES)* 7(3), 195–200.

Arbea, J.I., 2021. Checklist de Fauna Ibérica. Clase Collembola Lubbock, 1870 (Hexapoda) en la península ibérica, islas Baleares y Macaronesia (edición 2021). In: *Documentos Fauna Ibérica*, 16. Ramos MA, Sánchez Ruiz M, editors. Madrid: Museo Nacional de Ciencias Naturales, CSIC. 41 pp.

Arbea, J.I., Kahrarian, M., 2015a. The genus *Folsomides* Stach (Collembola, Isotomidae) in Kermanshah Province (W Iran) with the description of two new species. *Zootaxa* 3925(2), 281–290. <https://doi.org/10.11646/zootaxa.3925.2.9>

Arbea, J.I., Kahrarian, M., 2015b. Two new species and new data of Isotomidae Schaeffer, 1896 (Collembola: Entomobryomorpha) from Iran. *Arquivos Entomol6gicos* 14, 71–88.

Axelson (Linnaniemi), W.M., 1907. Die Apterygotenfauna Finlands. I. Allgemeiner teil. *Acta Societatis Scientiarum Fennicae* 34(7), 1–134.

Axelson, W.M., 1900. Vorläufige Mittheilung über einige neue Collembolen-Formen aus Finnland. *Meddelanden af Societatis pro Fauna et Flora Fennica* 26, 105–123. <https://doi.org/10.5962/bhl.part.26726>

Axelson, W.M., 1902. Diagnosen neuer Collembolen aus Finnland und angrenzenden Teilen des nordwestlichen Russlands. *Meddelanden of Societas pro Fauna et Flora Fennica* 28, 101–111.

Axelson, W.M., 1905. Einige neue Collembolen aus Finnland. *Zoologischer Anzeiger* 28(24/25), 788–794.

Babenko, A., Stebaeva, S., Turnbull, M.S., 2019. An updated checklist of Canadian and Alaskan Collembola. *Zootaxa* 4592(1), 1–125. <https://doi.org/10.11646/zootaxa.4592.1.1>

Bagnall, R.S., 1939. Notes on British Collembola. *Entomologist's Monthly Magazine* 7, 91–102 (IV) and 188–200 (V).

Bagnall, R.S., 1940. Notes on British Collembola. VI. *Entomologist's Monthly Magazine* 76, 97–102 (VI) and 163–174 (VII).

Bagnall, R.S., 1949. Notes on British Collembola. IX. *The Entomologist's monthly magazine* 85, 51–61.

Baijal, H.N., 1955. Entomological Survey of the Himalayas Part XI. -On Five New Species of Collembola. *Agra University Journal of Research (Science)* 4(2), 531–538.

Bakhshi, A., Shayanmehr, M., Mohammadi Sharif, M., Yahyapour, E., Kaprus', I., 2022. A faunistic study of springtails (Hexapoda, Collembola) from Hezarjirib forests (Neka, Mazandaran) with three new records of Iran. *Journal of Insect Biodiversity and Systematics* 8(3), 395–410. <https://doi.org/10.52547/jibs.8.3.395>

Balvasi, A., Chelav, H.S., Khashaveh, A., Shayanmehr, M. 2015. New report of Collembola (Hexapoda: Entognatha) from North of Iran (Larijan Region). *International Journal of Agriculture and Biosciences* 5(1), 24–28.

Baquero, E., Hamra-Kroua, S., Jordana, R., 2009. A new species of Entomobrya from Northern Algeria (Collembola: Entomobryidae). *Entomological News* 120(1), 65–75. <https://doi.org/10.3157/021.120.0113>

Baquero, E., Mandal, G., Jordana, R., 2015. Entomobryoidea (Collembola) from Himachal Pradesh (India) in the Himalayas. *Zootaxa* 4027(1), 1–41. <https://doi.org/10.11646/zootaxa.4027.1.1>

Baquero, E., Schulz, H-J., Moravvej, A., Jordana, R., 2008. On the identity and geographical distribution of *Entomobrya handschini* Stach, 1922 (Collembola, Entomobryidae). *Abhandlungen der Naturforschenden Gesellschaft zu Görlitz* 79(2), 257–269.

Bellinger P. F., Christiansen K. A. & Janssens F. 2024. Checklist of the Collembola of the World. Retrieved from: <http://www.collembola.org>.

Bonet, F., 1942. Notas sinonímicas sobre el orden Colembolos. *Ciencia* 3(2), 56–59.

Börner, C., 1901. Ueber ein neues Achorutiden genus *Willemia* sowie 4 weitere neue Collembolen formen derselben Familie. *Zoologischer Anzeiger* 24, 422–433.

Börner, C., 1903. Neue altweltliche Collembolen, nebst Bemerkungen zur Systematik der Isotomini und Entomobryinen. *Sitzungsberichten der Gesellschaft naturforschender Freunde zu Berlin* 129–182. <https://doi.org/10.5962/bhl.part.29866>

Börner, C., 1906. Das system der Collembolan, nebst beschreibungen neuer Collembolen des Hamburger Naturhistorischen Museums. *Mitteilungen aus dem Naturhistorischen Museum in Hamburg* 23, 147–188.

Börner, C., 1913. Die Familien der Collembolen. *Zoologischer Anzeiger* 41, 315–322.

Bourlet, C., 1839. Mémoires sur les Podures. *Mémoires de la Société Royale des Sciences de l'Agriculture et des Arts de Lille* 1, 377–417.

Brook, G., 1882. On a new Genus of Collembola (*Sinella*) allied to *Degeeria*, Nicolet. *Zoological Journal of the Linnean Society* 16(95), 541–545. <https://doi.org/10.1111/j.1096-3642.1882.tb02398.x>

Brown, J.M., 1926. On some Collembola from Mesopotamia. *Zoological Journal of the Linnean Society, London* 36(242), 201–218.

Carl, J., Lebedinsky, J., 1905. Materialien zur Höhlenfauna der Krim. *Zoologischer Anzeiger* 28(16–17), 562–565.

Cassagnau, P., Delamare Deboutteville, C., 1955. Mission Henri Coiffait (1951). 3. *Collemboles*. *Archives de Zoologie expérimentale et générale* 91, 365–395.

Christiansen, K., 1958. The Entomobryiform Male Genital Plate. *Proceedings of The Iowa Academy of Science* 65(20), 474–476.

Christiansen, K.A., 1957. The Collembola of Lebanon and Western Syria. Part II. Families Cyphoderidae and Oncopoduridae. *Psyche* 64(3), 77–89. <https://doi.org/10.1155/1957/32941>

Cox, P., 1982. The Collembola fauna of north and north western Iran. *Entomologist's Monthly Magazine* 118, 39–43.

Daghighi, E. 2012. Fauna of Collembola (Insecta: Apterygota) from Rasht and its regions. MSc thesis, Guilan, Vol. 1. University of Guilan, Iran, 97pp. [in Persian with English abstract].

Daghighi, E., Hajizadeh, J., Hosseini, R., Moravvej, A. 2013. Introduction of eighteen species of springtails (Arthropoda: Collembola) from Guilan province with three new records for Iran. *Entomofauna* 13, 177–184.

Dalla Torre, K.W. von, 1895. *Die Gattungen und Arten der Apterygogenea (Brauer)*. Sechshundvierzigstes Programm des K.K. Staats-Gymnasiums. Innsbruck: Wagner. 23 pp.

Dányi, L., Traser, G., 2008. An annotated checklist of the springtail fauna of Hungary (Hexapoda: Collembola). *Opuscula Zoologica, Budapest* 38, 3–82.

Darvish-Motevalli, M. 2016. Faunistic survey and succession of insects in a rice field from planting to harvesting time. MSc thesis, Sari, Iran: Sari Agricultural Science and Natural Resources University, 148 pp.

Deharveng, L., Christian, E., 1984. *Gnathofolsomia palpata* ng, n. sp., eine Isotomide mit abgewandelten Mundwerkzeugen aus österreichischen Höhlen (Insecta, Collembola). *Verhandlungen des Zoologisch-Botanischen Vereins in Wien* 122, 97–101.

Deharveng, L., Lek, S., 1993. Remarques sur la morphologie et la taxonomie du genre *Isotomurus* Börner, 1903 et description de deux espèces nouvelles de France (Collembola: Isotomidae). *Annales de la Société entomologique de France, New Series* 29, 245–259.

Denis, J.R., 1923. Notes zur les Aptérygotes. II. Sur la faune française des Aptérygotes. *Annales de la Société entomologique de France* 92, 237–246.

Denis, J.R., 1924. Sur les Collembolen du Muséum de Paris (1re partie). *Annales de la Société Entomologique de France* 93, 211–260.

Denis, J.R., 1931. Collembolen de Costa Rica avec une contribution au species de l'ordre. *Bollettino del Laboratorio di zoologia generale e agraria della R. Scuola superiore d'agricoltura in Portici* 25, 69–170.

Desor, E., 1841. Excursions et séjour de M. Agassiz sur la mer de glace du Lauteraar et du Finsteraar, en société de plusieurs naturalistes. *Bibliothèque Universelle de Genève, Nouvelle Serie* 32, 116–399.

Dunger, W.G., 1982. Collembolen (Insecta, Collembola) aus der Mongolischen Volksrepublik, II. Isotomidae. *Annales Historico-Naturales Musei Nationalis Hungarici* 74, 35–74.

Fabricius, J.C., 1775. *Systema entomologiae, sistens insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus*. Flensburgi and Lipsiae: Officina Libraria Kortii. 832 pp. <https://doi.org/10.5962/bhl.title.36510>

Falahati Hossein Abad, A., Potapov, M., Sarailoo, M.H., Shayan Mehr, M. 2013. New records of Isotomidae (Collembola) from Golestan province (Iran). *Munis Entomology and Zoology* 8, 236–238.

Falahati Hosseinabad, A., Sarailoo, M.S., Jafari, S.A., Alizadeh, S., 2011. New species report of family of Entomobryidae (Collembola) for Iranian fauna. *Journal of Novel researches on Plant Protection* 3(4), 315–328.

Fjellberg, A., 2007. *The Collembola of Fennoscandia and Denmark. Part II: Entomobryomorpha and Symphypleona*. Leiden, Boston, Köln: Brill.

Folsom, J.W., 1902. Collembola of the grave. *Psyche, Cambridge* 9(315), 363–367. <https://doi.org/10.1155/1902/42417>

- Frenzel, G., 1941. Eine neue Collembolenart: *Proisotoma marchica* nov. spec. *Märkische Tierwelt* 4, 321–324.
- Ghahramaninezhad, S., Shayanmehr, M., Yoosefi Lafooraki, E. 2013. Report of new Collembola (Arthropoda: Hexapoda) species from Kermanshah. *Journal of Plant Protection* 27, 136–138.
- Ghasemi Charati, M., Shayanmehr, M., Amiri Besheli, B., Gioias Cipola, N., 2022. Introduction to class of Collembola as soil mesofauna from Semeskandeh mixed forest (Hyrcanian region). *Ecology of Iranian Forests* 9(18), 115–126 (in Persian with English abstract).
- Ghasemi Cherati, M. 2017. The comparison fauna, diversity and density of Collembola in an agricultural ecosystem versus a forest ecosystem. MSc thesis, Sari, Iran: Sari Agricultural Science and Natural Resources University
- Ghavami, R. 2016. A Faunistic study on springtails (Hexapoda: Collembola) in Khorramabad county (Lorestan, Iran). MSc, Islamic Azad University, Arak Branch, Iran.
- Gisin, H., 1949. Notes sur les Collemboles avec description de quatorze espèces et d'un genre nouveaux. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 22(4), 386–409.
- Gisin, H., 1953. Notes sur les Collemboles, avec description de cinq espèces nouvelles découvertes dans le canton de Genève. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 26, 56–62.
- Gmelin, J.F., 1788. *Caroli A Linné, Systema Naturae. Tom. 1. Pars V.* Lipsiae, Beer, 2225–3020.
- Greenslade, P., 1992. The identity of Australian specimens recorded as *Lepidosinella armata* Handschin 1920 (Collembola: Entomobryidae) with a key to Australian *Sinella* and *Coecobrya*. *Journal of the Australian Entomological Society* 31, 327–330
- Gruia, M.M., Poliakov, P., Broza, M., 2000. Collembola of Northern Israel, II. *Contributions from the Biological Laboratory, Kyoto University* 29, 117–131.
- Handschin, E., 1924. Die Collembolenfauna des Schweizerischen Nationalparks. *Denkschriften der Schweizerischen naturforschenden gesellschaft* 60(2), I–VIII + 89–174.
- Hopkin, S.P., 1997. Biology of Springtails (Insecta: Collembola). Cambridge: Oxford University Press UK. 330 pp.
- Hosseini, F-S., Shayanmehr, M., Amiri-Besheli, B. 2016. Contribution to Collembola (Hexapoda: Entognatha) fauna from Golestan province, Iran. *Journal of Insect Biodiversity and Systematics* 2(3), 321–338.
- Jacquemart, S., 1974. Resultats de la mission Anthropologique Belge au Niger. Collemboles nouveau du Sahara. *Bulletin de l'Institut royal des Sciences naturelles de Belgique. Entomologie* 50(6), 1–46.
- Jordana, R., 2012. Capbryinae and Entomobryini. In: Dunger W & Burkhardt U, editors. *Synopses on Palearctic Collembola*. Görlitz: Senckenberg Museum of Natural History Görlitz. pp 1–390.

Jordana, R., Ardanaz, A., 1981. Contribución al conocimiento de los colémbolos de Navarra. *Xenyllodes monoculatus* n. sp., *Uzelia setifera clavata* n. ssp. y *Folsomia sexoculata trisetata* n. ssp. (Collembola) de Navarra. *Publicaciones de Biología de la Universidad de Navarra, Serie Zoológica* 6, 33–46.

Jordana, R., Baquero, E., 2010. A new species of *Entomobrya* from Iraq (Collembola, Entomobryidae). *Soil organisms* 82(3), 351–356.

Kahrarian, M. 2015. Study on population dynamics, biodiversity, fauna and habitat suitability modeling of springtails (Apterygota: Collembola) in Kermanshah province. PhD, Islamic Azad University, Arak, Iran Branch.

Kahrarian, M. 2019. The checklist of Collembola (Hexapoda, Arthropoda) from west of Iran. *Journal of Insect Biodiversity and Systematics* 5 (1), 33–46.

Kahrarian, M., Nikpy, A., Mohammadi Noor, L. 2012. Preliminary checklist of the Collembolan fauna in Kermanshah, Sahneh and Harsin counties (Kermanshah: Iran) with two new records for Iranian fauna. *Pakistan Entomologist* 34(1), 27–30.

Kahrarian, M., Vafaei-Shoushtari, R., Soleyman-Nejadian, E., Shayanmehr, M., Shams Esfandabad, B. 2015. New records of Isotomidae and Paronellidae for the Iranian fauna with an update Checklist of Entomobryomorpha fauna (Collembola) in Kermanshah province. *Journal of Entomological Research* 7(4), 55–68.

Kahrarian, M., Vafaei-Shoushtari, R., Soleymannezhadian, E., Shayan Mehr, M., Shams, B. 2014. A faunistic study on Entomobryidae (Collembola) in Kermanshah (Iran). *Natura Somogyiensis* 24, 17–24.

Lie-Pettersen, O.J., 1896. Norges Collembola, Fortagnelse over de i Norge hidtilte observerede Arten. *Bergens Museums Aarbog* 8, 3–26.

Lima, E.C.A. de, Zeppelini, D., 2021. First survey of Collembola (Hexapoda: Entognatha) fauna in soil of Archipelago Fernando de Noronha, Brazil. *Florida Entomologist* 98(1), 368–369. <https://doi.org/10.1653/024.098.0161>

Linnaeus, C., 1758. *Systema naturae*. 10th ed. Holmiae: Impensis Direct. Laurentii Salvii. Pp 608–609.

Lubbock, J., 1862. Notes on the Thysanura. Part II. *Transactions of the Linnean Society of London* 23, 589–601. <https://doi.org/10.1111/j.1096-3642.1860.tb00149.x>

Lubbock, J., 1868. Notes on the Thysanura.—Part III. *Transactions of the Linnean Society of London* 26(1), 295–304. <https://doi.org/10.1111/j.1096-3642.1968.tb00508.x>

Lubbock, J., 1870. Notes on the Thysanura. Part III. *Transactions of the Linnean Society of London* 26, 295–304. <https://doi.org/10.1111/j.1096-3642.1968.tb00508.x>

Lubbock, J., 1873. *Monograph of the Collembola and Thysanura*. Ray Society, London. 276 pp.

MacGillivray, A.G., 1896. The American species of *Isotoma*. *The Canadian Entomologist* 28(3), 47–58. <https://doi.org/10.4039/Ent2847-2>

- Mandal, G.P., 2018. Three new species of Collembola from India. *Records of the Zoological Survey of India* 118(2), 107–127.
- Mari Mutt, J.A., 1979. *Heteromurítrella*, a new tropical subgenus of *Heteromurus* with descriptions of two new species (Insecta: Collembola: Entomobryidae). *The Journal of Agriculture of the University of Puerto Rico* 63(2), 214–222. <https://doi.org/10.46429/jaupr.v63i2.10278>
- Mari Mutt, J.A., Stomp, N., 1980. Two new species of *Heteromurus* s. str. from Portugal and Romania (Insecta, Collembola, Entomobryidae). *Archives d' Institut Grand-Ducal de Luxembourg, Section des Sciences naturelles, physiques et mathématiques* 38, 183–189.
- Martynova, E., 1969. Springtails of the family Tomoceridae (Collembola) from the fauna of the USSR. *Entomological Review* 43, 299–314 [in Russian]
- Martynova, E.F., 1968a. Materials on fauna of springtails (Collembola) of middle Asia. *Izvestiya otdeleniya biologicheskikh nauk AN Tadzhikskoi SSR* 28, 32–46. (in Russian)
- Martynova, E.F., 1968b. New species of Springtails (Collembola) of the family Isotomidae from mountains regions of Middle Asia. *Zoologicheskii Zhurnal, Moscow* 47, 1556–1561. (in Russian)
- Martynova, E.F., 1971. New species of springtails (Collembola) from mountain and steppe regions of the USSR. *Revue d'Entomologie de l'USSR* 50(3), 598–611. (in Russian)
- Martynova, E.F., Chelnokov, V.G., Rasulova, Z.K., 1974. Species of the genus *Heteromurus* Wankel, 1860 (Collembola, Entomobryidae, s.l.) in the USSR fauna. *Vestnik Zoologii* 2, 66–71. (in Russian)
- Mateos, E., 2012. The European *Lepidocyrtus lanuginosus* group (Collembola: Entomobryidae), definition and description of a new species from Spain. *Zootaxa* 3570(1), 69–81. <https://doi.org/10.11646/zootaxa.3570.1.5>
- Mehrafrooz Mayvan, M., Greenslade, P., 2020. *Isotomurus graminis* Fjellberg (Collembola: Isotomidae) new for Iran, key to Iranian species in genus with comment on the validity of photographic records. *Zootaxa* 4838(3), 445–450. <https://doi.org/10.11646/zootaxa.4838.3.10>
- Mehrafrooz Mayvan, M., Sadeghi-Namaghi, H., Shayanmehr, M., Greenslade, P., 2022. Contribution to the knowledge of Entomobryomorpha (Hexapoda: Collembola) from Northeastern Iran with new records and a key to the species. *Journal of Insect Biodiversity and Systematics* 8(2), 151–174. <https://doi.org/10.52547/jibs.8.2.151>
- Moniez, R., 1899. Notes sur les Thysanoures—I: Espèces qui vivent aux Açores. *Revue Biologique du Nord de la France* 2, 24–31.
- Moradi, T., Vafaei-Shoushtari, R., Kahrarian, M., Mohseni-Amin, A. 2018. The study on Springtails in west part of Iran with new records for Iranian fauna. *Journal of Entomological Research* (10)3, 43-53.
- Moravvej, S.A. 2003. Biodiversity of Collembola of Tehran Region and Preliminary Observation on Several Species. MSc thesis, Tehran, Vol. 1. Tarbiat Modarres University, Iran, 135 pp. [in Persian with English abstract]

Moravvej, S.A., Potapov, M., Kamali, K., Hodjat, S.H. 2007. Isotomidae (Collembola of the Tehran region, Iran). *Zoology in the Middle East* 41, 117–118. <https://doi.org/10.1080/09397140.2007.10638238>

Müller, O.F., 1776. *Zoologiae Danicae prodromus, seu animalium Daniae et Norvegiae indigenarum characteres, nomina et synonyma imprimis popularium, Havniae XXXII*. Copenhagen: Hallageriis. 282 pp. <https://doi.org/10.5962/bhl.title.63795>

Negri, I., Pellecchia, M., Fanciulli, P.P., 2005. Two new species within the genus *Seira* Lubbock, 1869 from Morocco (Collembola, Entomobryidae). *Zootaxa* 840, 1–12. <https://doi.org/10.11646/zootaxa.840.1.1>

Nematollahi, M., Bagheri, M., Radwanski, J. 2009. New reports of Collembola for Iran with surveying of the importance in the greenhouses of Isfahan province, Iran. *Plant Protection Journal*, Islamic Azad University, Shiraz branch 3, 327–335. [in Persian with English abstract]

Neuherz, H., 1976. *Drepanosira hussi* sp. n.—eine neue Collembolenart (Insecta, Apterygota) aus dem Östlichen Hindukusch (Afghanistan). *Sitzungsberichte der Österreichische Akademie der Wissenschaften in Wien, Mathematisch-naturwissenschaftliche Klasse* 185, 1–4.

Nicolet, H., 1842. Recherches pour Servir à l'Histoire des Podurelles. *Nouveaux mémoires de la Société helvétique des sciences naturelles* 6, 1–88.

Nosek, J., 1964. Einige neue Collembolenarten aus Tadschikistan. *Zoologischer Anzeiger* 172(2), 140–146.

Packard, A.S., 1873. Synopsis of the Thysanura of Essex County, Mass., with descriptions of a few extralimital forms. *Fifth Annual Report of the Peabody Academy of Science* 1873, 23–51.

Paclt, J., 1944. Nomina nova in Collembola. *Entomologické listy (Folia Entomologica)* 7, 92.

Palissa, A., 1966. Zur Collembolen fauna Finnlands. *Notulae Entomologica* 46, 33–63.

Parona, C., 1888. Collembola e sui Tisanuri. *Annali del Museo civico di storia naturale di Genova* 6, 133–144.

Potapov, M., Deharveng, L., Janion-Scheepers, C., 2021. Taxonomy of the *Proisotoma* complex. VI. Rediscovery of the genus *Bagnallella* Salmon, 1951 and epitoky in *Bagnallelladavid* (Barra, 2001), comb. nov. from South Africa. *ZooKeys* 1072, 185–204. <https://doi.org/10.3897/zookeys.1072.71307>

Potapov, M., Kahrarian, M., Deharveng, L., Shayanmehr, M., 2015. Taxonomy of the *Proisotoma* complex. V. Sexually dimorphic *Ephemerotoma* gen. nov. (Collembola: Isotomidae). *Zootaxa* 4052(3), 345–358. <https://doi.org/10.11646/zootaxa.4052.3.4>

Potapov, M.B., Stebaeva, S.K., 1990. Species of the genus *Anurophorus* Nicolet, 1842 (Collembola: Isotomidae, Anurophorinae) in the fauna of the USSR. In: *Novye i maloizvestnye vidy fauny Sibiri (New or little known species in the fauna of Siberia)*. Novosibirsk: Nauka. 101 pp. (in Russian)

Qazi, F., Shayanmehr, M. 2014. Additional records for Iranian Collembola (Hexapoda: Entognatha) fauna from Tehran province. *Natura Somogyiensis* 25, 27–34.

- Qazi, F., Shayanmehr, M., 2016. A checklist of Collembola of Tehran, with some new records from Iran. *Journal of Entomological Society of Iran* 36(2), 121–136.
- Rabieh, M.M., Shayanmehr, M., Yoosefi Lafooraki, E., Noei, J., 2023. Three new records of Collembola (Hexapoda) from Birjand (South Khorasan, Iran). *Journal of Insect Biodiversity and Systematics* 9(3), 527–533.
- Ramezani, L., Mossadegh, M.S. 2017. The effect of cropping on diversity and density of springtails (Hexapoda: Collembola) in Khuzestan province, Southwest of Iran. *Journal of Entomological Research* 8(4), 51–57.
- Reuter, O.M., 1876. Catalogus præcursorius Poduridarum Fenniae. *Meddelanden Af Societas Pro Fauna Et Flora Fennica* 1, 78–86.
- Reuter, O.M., 1895. *Finlands Collembola och Thysanura*. Helsingfors: Apterygogenea Fennica. 203 pp.
- Rusek, J., 1981. Some Collembola from Iraq. *Vestnik Ceskoslovenske Spolecnosti Zoologicke* 45, 63–80.
- Salmon, JT., 1951. Keys and bibliography to the Collembola. *Zoology Publications from Victoria University College* 8, 1–82.
- Schäffer, C., 1896. Die Collembolen der Umgebung von Hamburg und benachbarter Gebiete. *Mitteilungen aus dem Naturhistorischen Museum in Hamburg* 13, 149–216.
- Schäffer, C., 1897. *Apterygoten*. Hamburg: Hamburger Magalhaensische Sammelreise. 48 pp.
- Schäffer, C., 1900. Ueber württembergische Collembola. *Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg* 56, 245–280.
- Schött, H., 1891. Beiträge zur Kenntniss kalifornischer Collembola. *Kongliga Svenska Vetenskaps-Akademiens Handlingar* 17, 1–25.
- Schött, H., 1893. Zur Systematik und Verbreitung palaearctischer Collembola. *Kongliga Svenska Vetenskapsakademiens Handlingar* 25, 1–100. <https://doi.org/10.5962/bhl.title.8645>
- Schött, H., 1896. North American Apterygogenea. *Proceedings of the California Academy of Sciences, Series 2* 6, 169–196.
- Schött, H., 1902. Études sur les Collembolés du Nord. *Bihang Svenska Vetenskaps-Akademiens Handlingar* 28(4), 1–48.
- Semati, F., Fakoorziba, M.R., Moemenbellah-Fard, M.D., Sarkari, B., Shayanmehr, M., 2023. A new record of springtails, *Drepanura pallens* Rusek, 1981 (Collembola: Entomobryidae) accessed by delusional parasitosis (DP) from south of Iran. *Journal of Asia-Pacific Biodiversity* 17(1), LicenseCC BY-NC-ND 4.0. <https://doi.org/10.1016/j.japb.2023.07.006>
- Shayanmehr, M., Marouf A., Yoosefi Lafooraki, E., Cipola, N., 2020a. *Seira atlantica* Negri, Pellecchia & Fanciulli, 2005: New record of Collembola (Entomobryidae: Seirinae) from alfalfa field in Iran. *Journal of Insect Biodiversity* 015(1), 1–5. <https://doi.org/10.12976/JIB/2020.15.1.1>.

Shayanmehr, M., Mirab-balou, M., Yoosefi Lafooraki, E., Kaprus', I., 2019. Introduction to Springtails (Arthropoda: Hexapoda) fauna from Ilam province (western Iran) with a new record from Sminthuridae. *Journal of Insect Biodiversity and Systematics* 5(4), 399–410. <https://doi.org/10.52547/jibs.5.4.399>

Shayanmehr, M., Yahyapour, E., Kahrarian, M., Yoosefi Lafooraki, E., 2013. An introduction to Iranian Collembola (Hexapoda): an update to the species list. *ZooKeys* 335, 69–83. <https://doi.org/10.3897/zookeys.335.5491>.

Shayanmehr, M., Yoosefi Lafooraki, E., Kahrarian, M., 2020b. A new updated checklist of Iranian Collembola (Arthropoda: Hexapoda). *Journal of Entomological Society of Iran* 39(4), 403–445. <https://doi.org/10.22117/jesi.2019.124366.1285>.

Shayanmehr, M., Yoosefi Lafooraki, E., Tully, T., D'Haese, C., Ghajar Sepanlou, M., 2024. New records of springtails (Hexapoda: Collembola) for Iran from the Bula Hyrcanian forest. *Journal of Insect Biodiversity and Systematics* 10(1), 31–42.

Shoebotham, J.M., 1917. Notes on Collembola—Part 4. The Classification of the Collembola; with a list of genera known to occur in the British Isles. *Annals and Magazine of Natural History, Series 8* 19(114), 425–436. <https://doi.org/10.1080/00222931709486959>

Shoeibi, M., Shayanmehr, M., Yoosefi Lafooraki, E., Mehrafrooz Mayvan, M. 2013. Faunistic survey of Entomobryidae family (Collembola; Hexapoda) in Iran. *Bulltin information Society of Lepidoptera of Romania* 23, 38-42.

Stach, J., 1922. Apterygoten aus dem nordwestlichen Ungarn. *Annales Musei Nationalis Hungarici* 19, 1–75.

Stach, J., 1947. *The Apterygotan fauna of Poland in relation to the world-fauna of this group of insects. Family: Isotomidae*. Kraków: Acta Monographica Musei Historiae Naturalis. 488 pp.

Stach, J., 1960. Materialien zur Kenntnis der Collembolen Fauna Afghanistans. *Acta Zoologica Cracoviensia* 5, 507–581.

Stach, J., 1963. *The apterygotan fauna of Poland in relation to the world fauna of this group of insects. Tribe: Entomobryini*. Kraków: Polska Akademia Nauk, Zakład Zoologii Systematycznej, Państwowe Wydawnictwo Naukowe. 126pp.

Stach, J., 1965. On some Collembola of North Vietnam. *Acta zoologica cracoviensia* 10(4), 345–371.

Templeton, R., Westwood, J.O., 1836. Thysanura Hibernica, or Descriptions of such Species of Spring-tailed Insects (*Podura* and *Lepisma*, Linn.) as have been observed in Ireland. *The Transactions of the Entomological Society of London* 1, 89–98.

Thibaud, J.M., 2017. Catalogue des collemboles de France. *Zoosystema* 39(3), 297–436. <https://doi.org/10.5252/z2017n3a1>

Tullberg, T., 1876. Collembola borealia (Nordiska Collembola). *Ofversigt af Kongliga Vetenskaps-Akademiens Forhandlingar* 33, 23–42.

Tullberg, T., 1877. *Collembola borealia*. - Nordiska Collembola., Ofversigt af Kongl. Vetenskaps Akademiens-Förhandlingar 5, 23–42.

Tullberg, T.F., 1871. Forteckning ofver Svenska Podurider. *Kongliga Svenska Vetenskaps-akademiens handlingar* 28(1), 143–155.

Uzel, J., 1890. *Thysanura Bohemiae*. *Sitzungsberichte der Königlich böhmischen Gesellschaft der Wissenschaften* 2, 3–82.

Vahedi Moghadam, S., Shayanmehr, M., Mohamadi Sharif, M., 2022. Springtails fauna (Hexapoda, Collembola) from different ecosystems of Behshahr and suburb (Mazandaran) with new species records. *Journal of Insect Biodiversity and Systematics* 8(4), 581–593. <https://doi.org/10.52547/jibs.8.4.581>

Wahlgren, E., 1899. Über die von der schwedischen Polarexpedition 1898 gesammelten Collembolen. *Öfversigt af Konglig Vetenskaps-Akademiens Förhandlingar* 56, 335–340.

Wankel, H., 1860. Beiträge zur fauna der Mäharichen Höhlen. *Lotos, Prague* 10, 201–206.

Willem, W., 1902. Note préliminaire sur les Collemboles des grottes de Han et de Rochefort. *Annales de la Société entomologique de Belgique* 46, 275–283.

Winkler, D., Mateos, E., Traser, G., Lakatos, F., Tóth, V. 2020. New Insight into the Systematics of European *Lepidocyrtus* (Collembola: Entomobryidae) Using Molecular and Morphological Data. *Insects* 11(5), 302. <https://doi:10.3390/insects11050302>

Yadav, R.S., Prasad, S., Kumar, D., 2018. Diversity of epigeic Collembola of Bibhutibhusan Wildlife Sanctuary, North 24 Parganas, West Bengal (India). *International Journal of Fauna and Biological Studies* 5(2), 17–21

Yahyapour, E., Shayanmehr, M. 2011. First report of two genus and five species of Collembola (Hexapoda: Entognatha) from Iran. *Plant Protection Journal* 3(1), 37-51.

Yahyapour, E., Vafaei-Shoushtari, R., Shayanmehr, M. Arbea, J. 2018. A survey on Entomobryomorpha (Collembola) fauna in northern forests of Iran. *Journal of Insect Biodiversity and Systematics*, 4(4), 307–316.

Yahyapour, E., Vafaei-Shoushtari, R., Shayanmehr, M., 2019. Study on the Collembola Fauna of Mazandaran (Iran) and new records from the forests of this province. *Journal of Entomological Research* 10(4), 103–113.

Yahyapour, E., Vafaei-Shoushtari, R., Shayanmehr, M., 2020. First report of the genus *Aethiopella* Handschin, 1942 and species *Folsomia inoculata* Stach, 1947 (Hexapoda Collembola) from Iran. *Acta entomologica serbica* 25(2), 1–10. <https://doi.org/10.5281/zenodo.4028689>

Yahyapour, E., Vafaei-Shoushtari, R., Shayanmehr, M., Arbea, J., 2021. New records of springtails (Hexapoda: Collembola) for Iran from Mazandaran forests. *Journal of Insect Biodiversity and Systematics* 7(3), 263–276. <https://doi.org/10.52547/jibs.7.3.263>

- Yahyapour, E., Vafaei-Shoushtari, R., Shayanmehr, M., Arbea, J., 2018. A survey on Entomobryomorpha (Collembola) fauna in northern forests of Iran. *Journal of Insect Biodiversity and Systematics* 4(4), 307–316. <https://doi.org/10.52547/jibs.4.4.307>
- Yoosefi Lafooraki, E. 2014. Morphological and Biological identification of Collembola (Hexapoda) in Mazandaran province. MSc thesis, Sari, Iran: Sari Agricultural Science and Natural Resources University, 132 pp.
- Yoosefi Lafooraki, E., Hajizadeh, J., Shayanmehr, M., Hosseini, R., 2020. Isotomidae (Collembola) from northern Iran with description of a new species of *Isotomodes* Linnaniemi. *Journal of Asia-Pacific Biodiversity* 13, 545–553. <https://doi.org/10.1016/j.japb.2020.08.012>
- Yoosefi Lafooraki, E., Hajizadeh, J., Shayanmehr, M., Hosseini, R., 2019. First report of *Anurophorus silvaticus* (Collembola: Isotomidae) for Iran. *Iranian Journal of Animal Biosystematics (IJAB)* 15(2), 171–174. <https://doi.org/10.22067/ijab.v15i2.82180>
- Yoosefi Lafooraki, E., Shayanmehr, M. 2014. A survey on Entomobryomorpha (Collembola, Hexapoda) fauna in Northern Iran with an identification key. *Iranian Journal of Animal Biosystematics* 10 (2), 101–117.
- Yoosefi Lafooraki, E., Shayanmehr, M., Qazi, F. 2014. First report of *Pseudosinella baghdadica* (Collembola: Entomobryidae) from Iran. 21st Iranian Plant Protection Congress, 23-26 August 2014, Urmia University, Urmia, Iran.
- Yoshii, R., Suhardjono, Y.R., 1989. Notes on the Collembolan Fauna of Indonesia and its vicinities. I. Miscellaneous Notes, with special references to Seirini and Lepidocyrtini. *Acta Zoologica Asiae Orientalis* 1, 23–90.
- Yosii, R., 1955. Meeresinsekten der Tokara Inseln. VI. Collembolen nebst Beschreibungen terrestrischer Formen. *Publications of the Seto Marine Biological Laboratory* IV (2-3), 1955-05-30. (Article 34), 379–401.
- Yosii, R., 1956. Höhlencollembolen Japans II. *Japanese Journal of Zoology* 11(5), 609–627.
- Yosii, R., 1961. On some Collembola from Thailand. *Nature and Life in Southeast Asia* 1, 171–201.
- Yosii, R., 1963. On some Collembola of Hindukush, with notes on *Isotoma* Bourlet and its allies. In: *Results of the Kyoto University Scientific Expedition to the Karakoam and Hindukush, 1955, IV Insect Fauna of Afghanistan and Hindukush. Art. I.* Kyoto: Kyoto University. 43 pp.
- Yosii, R., 1966. On some Collembola of Afghanistan, India and Ceylon, collected by the Kuphe-Expedition, 1960. *Results of the Kyoto University Scientific Expedition to the Karakoram and Hindukush* 8, 333–405.
- Yu, D., Ding, Y., Tihelka, E., Cai, Ch., Hu, F., Liu, M., Zhang, F., 2022. Phylogenomics of Elongate-Bodied Springtails Reveals Independent Transitions from Aboveground to Belowground Habitats in Deep Time. *Systematic Biology* 71(5), 1023–1031, <https://doi.org/10.1093/sysbio/syac024>

Zamani Khormandichali, F., Shayanmehr, M., Yoosefi Lafooraki, E., Mohammadi Sharif, M., 2024. Springtails (Hexapoda, Collembola) of Savadkooh (Mazandaran) with the description of a new species of *Isotomurus* Börner (Entomobryomorpha, Isotomidae). *Journal of Insect Biodiversity and Systematics* 10(1), 59–72. <https://doi.org/10.61186/jibs.10.1.59>

Zamani, N. 2016. Investigation of hexapoda fauna in Rapeseed (*Brassica napus*: Brassicaceae) and introduction to pests and their natural enemies in Sari (Mazandaran: Iran). MSc thesis, Sari, Iran: Sari Agricultural Science and Natural Resources University.

Zhang, F., 2015. Some *Willowsia* from Nepal and Vietnam (Collembola: Entomobryidae) and description of onr new species. *Zootaxa* 3905(4), 489–499. <https://doi.org/10.11646/zootaxa.3905.4.3>